

The Role of Cystatin C in the Diagnosis of Renal Dysfunction in Patients with Coronary Heart Disease

M. E. Rakhimova, A. G. Gadaev

Tashkent Medical Academy, Tashkent, Uzbekistan

Abstract Diagnosis and treatment of chronic coronary heart disease (CHD) is well developed, but there are often difficulties in managing patients with comorbid diseases. A very common chronic kidney disease (CKD) occupies a special place among comorbidities. The cause of death in patients with kidney dysfunction is significantly more often CHD than end-stage renal disease (ESRD). In recent years, cystatin C has been considered as an early marker for assessing the functional state of the kidneys and cardiovascular risk, the level of which, unlike creatinine, does not depend on gender, age, race, and muscle mass.

Keywords Coronary heart disease, Cystatin C, Functional state of the kidneys

1. Introduction

Cardiovascular illness (CVD) is, without special case, one of the driving causes of passing and incapacity in populaces of all nations. Agreeing to master reports of the World Wellbeing Organization (WHO), affirmed by measurable investigation, atherosclerotic cardiovascular malady has been a driving cause of dismalness and mortality within the world populace (accounting for 55% of the full populace) for numerous a long time. is the most cause of More than two-thirds of these are ischemic heart illness (CHD). In 2012, 17.5 million individuals kicked the bucket from cardiovascular infection, bookkeeping for 31% of all passings around the world. Of these, 7.4 million passed on from coronary course illness and 6.7 million from stroke [11,12].

Populace maturing and way of life changes are expanding the predominance of cardiovascular malady in creating nations (up to 75% of cardiovascular infection passings happen in moo- and middle-income nations), Agreeing to WHO, a undermining upward slant in mortality has been watched.

An evaluated 22 to 25 million individuals will pass on from coronary course infection in 2025, about half of whom will kick the bucket from coronary supply route illness. It has been concluded that humankind is on the brink of a modern non-infectious widespread of the third thousand years [12].

The predominance of coronary supply route illness is around 10% of the grown-up populace in created nations and roughly 20% within the elderly and the elderly. As it were 40-50% of her patients are mindful of their condition. The remaining 50-60% stay undetected [3].

2. The Main Findings and Results

Based on numerous clinical, research facility, and epidemiological ponders, the improvement of atherosclerosis, counting coronary course malady, is associated with way of life, the nearness of certain metabolic highlights, malady or neurotic conditions. have been illustrated and together these are characterized as chance components. RF) for coronary course illness. The foremost critical of them are:

Smoking, diabetes mellitus (DM), blood vessel hypertension (Ok), weight, nearness of coronary course illness in a relative, and decreased physical action. The chances of creating coronary supply route infection increment when two, three, or more of the hazard components recorded combine, particularly with a inactive way of life [11]. In expansion to the conventional chance variables for CAD (DM, Ok), renal brokenness is related with expanded irritation, thoughtful movement, endothelial brokenness, atherosclerosis, plaque defenselessness, vascular calcification and frailty. Expanded thoughtful adrenal movement is common in renal brokenness (RD) and is related with an expanded hazard of movement to cardiovascular and renal infection. Renal brokenness is common in patients with hypertension, coronary heart infection, particularly diabetes and incessant heart disappointment (CHF). At the same time, renal brokenness is also an critical free hazard figure for the improvement of cardiovascular complications such as myocardial dead tissue (MI), heart disappointment, and lethal arrhythmias [4].

The kidney and its glomeruli are portion of the body's microcirculatory framework and impact the improvement of cardiovascular pathologies, whereas the kidney itself is additionally included within the neurotic forms of different cardiovascular infections. In later decades, the number of DP patients has expanded around the world. Concurring to

different population-based registries and thinks about (e.g., NHFES III, Okinawa Ponder, PREVEND, Swiss SAPALDIA study), the predominance of kidney illness is 10-13% within the Joined together States, Europe and Japan, with indeed higher levels at tall elevations. has come to - Chance Gather. At the same time, the increment in renal infection predominance is basically related with relentlessly expanding predominance of cardiovascular illness, sort 2 diabetes and weight [7].

The affiliation between renal brokenness and expanded frequency of cardiovascular disease and its complications within the common populace was to begin with examined within the Framingham Heart Ponder. The consider included 6,223 members (cruel age 54 a long time, 54male) and was taken after for 15 a long time. Serum creatinine was chosen as an marker of diminished renal work.

1.5 to 3.0 mg/dl for men and 1.4 to 3.0 mg/dl for ladies. An upper constrain of creatinine concentration was indicated to prohibit cases of extreme renal disappointment. Since the begin of the perception period, renal work compounded in 8.7% of men and 8.0% of ladies. At the conclusion of 15 a long time of follow-up, diminished renal function was not associated with an expanded chance of cardiovascular occasions in either men or ladies. Be that as it may, in men, hypercreatinineemia is related with an expanded hazard of all-cause mortality. The creators of this think about were compelled to report negative comes about, concluding that the expanded risk of cardiovascular complications is likely due to related cardiovascular chance components, instead of diminished renal work. [3].

A Japanese think about of 2,634 members matured 40 a long time or more seasoned and enduring 12 a long time (Hisagama think about) found a clear relationship between diminished GFR and the frequency of coronary course infection in men and the rate of coronary artery illness. critical pertinence was found. Ischemic stroke in ladies [3].

In spite of the fact that the conclusion and treatment of coronary supply route malady are well created, administration of patients with comorbidities is regularly troublesome. Constant kidney illness (CKD), which is exceptionally common, is considered a major hazard calculate for cardiovascular malady and possesses a extraordinary put among comorbidities. Coronary course infection is altogether more common than end-stage renal malady in DP patients [1]. Information from the EUROPA think about appear a tall frequency of PD in patients with coronary course malady. With steady coronary course illness, he had a GFR 50%) in 53% of patients with direct and extreme DP and 28% of patients with mellow or typical renal work [5].

Numerous huge worldwide thinks about have affirmed the tall predominance of CKD, comparable to that of coronary course illness and diabetes. Comes about from a CKD (Chronic Kidney Malady Forecast Consortium) prognostic think about appeared critical and free associations between GFR and albuminuria with all-cause and cardiovascular mortality, movement of CKD and hazard of intense kidney harm. [2].

An affiliation between renal brokenness and coronary atherosclerosis was affirmed at dissection.

For GFR ≥ 60 , 45-59, 30-44, and <30 ml/min/1.73 m², serious coronary atherosclerosis was recognized in 34, 42, 52, and 53% of cases . Within the Mettle think about, the nearness of persistent kidney infection was related with a 1.5-fold expanded chance of heart assault, passing and modern heart disappointment cases in patients with steady coronary course illness. The 2-year survival rate after intense MI was 57% for CKD stages 1-2 and 30% for stages 4-5. The chance of having a heart assault inside two a long time was higher for those with incessant kidney infection than for those with diabetes. In this way, a bidirectional relationship between CAD and DP is famous for expanded illness rate and more awful guess for CAD with more extreme CKD [1]. Diminished GFR as a hazard calculate for antagonistic results in people with basic cardiovascular infection. In one of the biggest randomized planned Trust ponders (Cardiac Results and Avoidance Appraisal), we to begin with compared the angiotensin-converting protein (Expert) inhibitor ramipril and cancer prevention agents on the rate of heart assault, stroke, and cardiovascular passing in 9297 a long time. The impact of treatment with the medicate vitamin E was examined. Tolerance. All subjects within the ponder had a place to the high-risk gather for creating cardiovascular occasions. Besides, the recurrence of essential cardiovascular infection (cardiovascular passing, intense myocardial infarction, stroke) was higher within the gather of patients with serum creatinine levels of 1.4–2.3 mg/dl (n=980) and within the bunch of ordinary working patients tried. Assessed. pointer. Kidney (n=8307). The same investigation was performed after calculating the GFR for bunches of individuals with values of this parameter underneath 65 ml/min (n=3394) and over 65 ml/min (n=5888). Indeed essentially comparing patients for creatinine or his GFR, more than one-third of patients (freely of other hazard variables and microalbuminuria) have a critical affect on essential cardiovascular results. It was shown that he had a few degree of renal disappointment. Cardiovascular mortality within the quiet bunch with marginally lifted serum creatinine was 22.2% vs. 15.1% [3]. This (autonomously of other hazard variables and microalbuminuria) had a noteworthy affect on major cardiovascular results. Cardiovascular mortality within the understanding bunch with marginally hoisted serum creatinine was 22.2% vs. 15.1% [3]. This (freely of other chance variables and microalbuminuria) had a critical affect on major cardiovascular results. Cardiovascular mortality within the understanding gather with somewhat raised serum creatinine was 22.2% compared to 15.1% [3].

Hillege HZ et al. Assessing different hemodynamic and serological variables as prognostic variables in patients with serious but steady heart disappointment, the most noteworthy indicator of all-cause mortality in this quiet category was the degree of GFR decay. Something startling turned out.

The lower the esteem, the higher the hazard of passing. These creators found that moo GFR values in patients with

AMI were prescient of the improvement of heart disappointment (counting those with cardiogenic stun and intense renal disappointment) [3].

As of late, cystatin C has been considered as a surrogate marker for evaluating renal utilitarian status and cardiovascular hazard. Cystatin C levels, not at all like creatinine, are not subordinate on muscle mass, permitting a more precise assurance of her GFR in individuals with the taking after illnesses:

Unusual physical make-up, need or abundance muscle mass. The weight of children, the elderly, diabetics, weight, and pregnant ladies. When the kidney is included in obsessive forms, cystatin C filtration within the kidney is impeded, driving to increased cystatin C substance within the blood. [13]. His estimation of GFR utilizing the CKD-EPI equation based on cystatin C or cystatin/creatinine predicts CAD seriousness more precisely than other equations (language structure ≥ 23) [26]. His GFR estimation utilizing the CKD-EPI equation anticipated coronary intercession results way better than the Cockcroft-Gault and MDRD equations [19].

Amassed involvement with expansive eyewitnesses proposes that GFR evaluation strategies based on estimation of cystatin C in blood have higher affectability and specificity than estimations of serum creatinine in people matured 16 a long time and more seasoned. It is appeared [16,17, 20].

At first tall at birth, cystatin C levels decrease within the to begin with year of life, stay stable until age 50, and after that increment [14].

Afshinnia F. And the co-authors found that cystatin C was the foremost delicate marker of early renal harm within the consider and a prognostic figure for renovascular illness in patients with fundamental hypertension, preeclampsia, and atherosclerotic changes. On the other hand, we considered that the seriousness of renal dysfunction was related with diastolic brokenness. Changes in left ventricular myocardium status and cleared out ventricular shape in numerous bunches. [5].

In patients with ischemic heart illness, serum cystatin C levels were straightly connected with systolic blood weight (SBP), indeed with ordinary renal work [23].

The creators too point out that the estimation of cystatin C in blood isn't as it were more precise, but too an early pointer of diminished kidney filtration. This characterization of circulating cystatin C as a marker of GFR decay is due to the reality that its concentration is free of sex and age (but for children beneath 1 year of age), but the protein is autonomous of skeletal muscle digestion system. It is ingested and not excreted into the renal tubules. Besides, raised blood cystatin C isn't only a indicator of unfavorable occasions such as renal harm, but also the risk of cardiovascular complications, for illustration in patients with CHF [6,8,21,24].

Numerous thinks about have appeared that lifted cystatin levels are related with an expanded hazard of mortality and cardiovascular occasions such as myocardial dead tissue, ischemic stroke, heart disappointment, fringe supply route

infection and metabolic disorder. In any case, most considers found cystatin to be prevalent to creatinine in exactness as a marker in these cases [9,10,15,18].

The increased cardiovascular chance related with expanded serum cystatin C is especially characteristic of the elderly [25], who commonly experience yearly decreases in her GFR. In addition, such a diminish in GFR is an autonomous chance calculate for expanded mortality [22]. Since the affiliation between serum cystatin C levels and long-term cardiovascular malady results is more grounded than the affiliation of such results with diminished GFR, raised cystatin C levels are related with cardiac It has been recommended that it may be related with expanded vascular mortality. [11].

In common, lifted circulating cystatin levels are related with expanded hazard of cardiovascular illness, hypertension, dyslipidemia, and mortality. Cystatin C levels are much less influenced by sex, age, race, muscle mass and weight than are creatinine levels. Various planned considers have appeared that a individual with elevated cystatin levels has an expanded hazard of creating her CVD and CKD in different clinical scenarios. Of specific esteem in recognizing those at high hazard for cardiovascular occasions, among those with typical GFR and creatinine and hence misclassified as moo chance for cardiovascular malady and persistent kidney malady, is the cystatin Chas It could be a level estimation.

3. Conclusions

Hoisted cystatin C levels are subsequently an pointer of preclinical renal disease related with cardiovascular chance and a indicator of its adverse complications. In common, cystatin C gets to be a reliable indicator of preclinical basic changes within the cardiovascular framework and serves as a more precise degree of renal work than serum creatinine or his GFR. Moreover, testing cystatin C levels gives more comprehensive prognostic data than other markers of renal work. Estimation of cystatin C alone or in combination with creatinine has the potential to more precisely degree GFR and evaluate hazard of passing and renal occasions in patients with coronary course illness.

REFERENCES

- [1] Belyalov F.I. Ischemic heart disease and impaired renal function. Rational pharmacotherapy in cardiology 2017; 13(3): 409-415.
- [2] Breidthardt, T. Diagnostic and prognostic value of cystatin C in acute heart failure / T. Breidthardt, Z. Sabti, R. Ziller, F. Rassouli, R. Twerenbold, N. Kozhuharov, E. Gayat, S. Shrestha, S. Barata, P. Badertscher, J. Boeddinghaus, T. Nestelberger, C. Mueller // Clin Biochem. - 2017. - Vol. 50, № 18. -P. 1007-1013.
- [3] Correa, S. Cystatin C for risk stratification in patients after an acute coronary syndrome / S. Correa, D.A. Morrow, E.

- Braunwald, R.Y. Davies, E.L. Goodrich, S.A. Murphy, C.P. Cannon, M.L. O'Donoghue // *J Am Heart Assoc.* - 2018. - Vol. 7, № 20.
- [4] Delanaye P, Cavalier E, Krzesinski JM et al Cystatin C, renal function, and cardiovascular risk // *Ann. Intern. Med.* 2008, 148 (4): 323.
- [5] Deng Y., Wang L., Hou Y. et al. The influence of glycemic status on the performance of cystatin C for acute kidney injury detection in the critically ill // *Ren. Fail.* - 2019. -Vol. 41 (1)—P. 139–149.
- [6] Deo R, Fyr CL, Fried LF, et al.. "Kidney dysfunction and fatal cardiovascular disease--an association independent of atherosclerotic events: results from the Health, Aging, and Body Composition (Health ABC) study". *Am. Heart J.* 2008. 155 (1): 62–8.
- [7] Eckel, R.H. 2013 AHA/ACC Guideline on Lifestyle Management to Reduce Cardiovascular Risk [Text] / R.H. Eckel, J.M. Jakicic, J.D. Ard, N.H. Miller, V.S. Hubbard, C.A. Nonas, J.M. de Jesus, F.M. Sacks, I.-M. Lee, Jr.S.C. Smith, A.H. Lichtenstein, L.P. Svetkey, C.M. Loria, T.W. Wadden, B.E. Millen, S.Z. Yanovski // *Journal of the American College of Cardiology.* – 2013, doi: 10.1016/j.jacc.2013.11.003.
- [8] Elsayed, M.S. Serum cystatin C as an indicator for early detection of diabetic nephropathy in type 2 diabetes mellitus / M.S. Elsayed, A. El Badawy, A. Ahmed, R. Omar, A. Mohamed // *Diabetes Metab Syndr.* - 2019. - Vol. 13, № 1. -P. 374-381.
- [9] Gadaev A.G., Rakhimova M.E., Turgunova M.U. The role of cystatin c in assessing the functional state of the kidneys in patients with coronary heart disease *Therapeutic Bulletin of Uzbekistan.* 2022. No. 4. C-104-110.e.
- [10] Glavnova O.B., Yarmolinskaya M.I., Suslova S.V., Borovik N.V. Possibilities of using cystatin C in the diagnosis of various diseases // *Journal of Obstetrics and Women's Diseases.* - 2018. - T. 67. - No. 4. - S. 40–47. doi: 10.17816/JOWD67440-47.
- [11] Kim S., Hwang S., Jang H. R. et al. Creatinine- and cystatin C-based estimated glomerular filtration rate slopes for the prediction of kidney outcome: a comparative retrospective study // *BMC Nephrol.* -2019. -Vol. 20 (1) - P. 214.
- [12] Liu, X. Is the chronic kidney disease epidemiology collaboration creatinine-cystatin C equation useful for glomerular filtration rate estimation in the elderly / X. Liu, H. Ma, H. Huang [et al] // *Clin Interv Aging.* - 2013. - № 8. -P. 1387-1391.
- [13] Luis-Lima, S. An overview of errors and flaws of estimated GFR versus true GFR in patients with diabetes mellitus / S. Luis-Lima, E. Porrini // *Nephron.* - 2017. - Vol. 136, № 4. - P. 287-291.
- [14] Osaki T., Satoh M., Tanaka F. et al. The Value of a Cystatin C-based Estimated Glomerular Filtration Rate for Cardiovascular Assessment in a General Japanese Population: Results from the Iwate Tohoku Medical Megabank Project // *J. Epidemiol.* - 2019. -May 25. doi: 10.2188/jea.JE20180274. [Epub ahead of print].
- [15] Parsh J., Seth M., Aronow H., et al. Choice of Estimated Glomerular Filtration Rate Equation Impacts Drug-Dosing Recommendations and Risk Stratification in Patients With Chronic Kidney Disease Undergoing PCI. *J Am Coll Cardiol.* 2015; 65(25): 2714-23.
- [16] Porrini, E. Remuzzi estimated GFR: time for a critical appraisal / E. Porrini, P. Ruggenenti, S. Luis-Lima, F. Carrara, A. Jiménez, A.P.J. de Vries, A. Torres, F. Gaspari // *GNat Rev Nephrol.* - 2018.
- [17] Ravn, B. Superiority of serum cystatin C over creatinine in prediction of long-term prognosis at discharge from ICU / B. Ravn, J.R. Prowle, J. Mårtensson, C.R. Martling, M. Bell // *Crit Care Med.* - 2017. -Vol. 45, № 9. - P. 932-940.
- [18] Scheen A.J. Pharmacokinetic considerations for the treatment of diabetes in patients with chronic kidney disease. *Expert Opin Drug Metab Toxicol* 2013; 9 (5): 529-550.
- [19] Shen, G. Increased cystatin C level in ST-elevation myocardial infarction predisposes the prognosis of angioplasty / G. Shen, H. Zhu, H. Ding, C. Sun, K. Zhou, Y. Fan, T. Li, M. Men, Y. Chen, Q. Lu, A. Ma // *Am J. Med Sci.* - 2018. - Vol. 355, № 6. - P. 530-536.
- [20] Shlipak MG, Katz R, Kestenbaum B, et al.: Rate of kidney function decline in older adults: a comparison using creatinine and cystatin C. *Am J Nephrol* 2009; 30: 171–8.
- [21] Waldeyer C., Karakas M., Scheurle C., et al. The predictive value of different equations for estimation of glomerular filtration rate in patients with coronary artery disease — Results from the AtheroGene study. *Int J Card.* 2016; 221: 908-13.