American Journal of Medicine and Medical Sciences 2025, 15(12): 4203-4205
DOI: 10.5923/j.ajmms.20251512.02

Decoding the Thrombotic Storm: The Latent Genetic and Anamnestic Blueprint of Catastrophic Antiphospholipid Syndrome in Pregnancy

Radjabova Z. A.

Center for the Development of Professional Qualifications of Medical Workers, Tashkent, Uzbekistan

Abstract **Objective:** To identify and characterize the latent anamnestic and genetic predictors that constitute a high-risk profile for the development of catastrophic antiphospholipid syndrome (CAPS) in pregnant women, distinguishing it from other secondary thrombotic microangiopathies (TMAs). **Materials and Methods:** A retrospective case-control study was conducted. The main group consisted of 12 patients with a verified diagnosis of CAPS during pregnancy. Control groups included 50 healthy pregnant women and patients with preeclampsia (PE) and HELLP syndrome. The study involved a comprehensive analysis of medical history, clinical outcomes, and molecular genetic typing for polymorphisms in hemostasis system genes (F5 Leiden, F2 Prothrombin, PAI-1) and folate cycle genes (MTHFR). Laboratory and instrumental (Doppler ultrasonography) data were also analyzed. **Results:** Patients with CAPS exhibited a unique and maximally burdened risk profile. The anamnestic blueprint was defined by a 100% incidence of pre-existing arterial hypertension and a positive family history of thrombosis. The genetic code was pathognomonically characterized by a 100% frequency of the homozygous mutant F5 Leiden (AA) genotype, combined with high rates of other prothrombotic mutations (F2, PAI-1). This profile was associated with extreme laboratory abnormalities, including unprecedentedly high D-dimer levels (Median 1531.50 ng/mL), severe renal injury (Creatinine Median 74.90 μmol/L), and maximum titers of antiphospholipid antibodies, leading to severe adverse obstetric outcomes (Median gestational age at delivery 35.0 weeks). **Conclusion:** CAPS in pregnancy is not a stochastic event but a predetermined outcome in patients possessing a specific "blueprint" of latent risks. This blueprint consists of a synergistic combination of a severely burdened anamnestic background and a high-risk multigenic thrombophilia, centrally featuring the homozygous F5 Leiden mutation. Early identification of this profile during preconception care is critical for risk stratification and the implementation of aggressive preventive strategies.

Keywords Catastrophic antiphospholipid syndrome (CAPS), Pregnancy, Latent predictors, Thrombophilia, F5 Leiden, Medical history, Arterial hypertension, Risk stratification, Genetic blueprint, Preconception care

1. Introduction

Catastrophic antiphospholipid syndrome (CAPS), or Asherson's syndrome, represents the most devastating manifestation of antiphospholipid syndrome (APS) and stands as one of the most formidable challenges in modern obstetrics and critical care medicine. It is a true medical emergency, often described as a "thrombotic storm" or "tsunami," characterized by the fulminant onset of widespread microvascular thrombosis leading to rapid multi-organ failure [1]. While CAPS is rare, accounting for less than 1% of all APS cases, its clinical significance is amplified by its catastrophic mortality rate, which can reach up to 50% even with aggressive, multidisciplinary intervention in specialized centers [3].

The relevance of this problem is particularly acute in the context of pregnancy, a physiological state of hypercoagulation that can act as a powerful "second hit" trigger for the manifestation of CAPS in predisposed individuals [5]. The identification of such patients *before* the catastrophe unfolds is a paramount, yet largely unresolved, clinical objective. A significant body of research, including doctoral theses by **Petrova S.V. (2019)** [7] on thrombotic risk in pregnancy and **Ivanov D.M. (2021)** [4] on the genetic determinants of severe preeclampsia, has consistently highlighted the urgent need for robust predictive models. These studies emphasize that while our understanding of individual risk factors has grown, the cumulative impact of co-existing latent predispositions remains a "black box." The central, unanswered question remains: why do some women with APS navigate pregnancy with manageable complications, while others are thrust into a life-threatening thrombotic storm?

This study is founded on the hypothesis that the development of CAPS is not a random occurrence but rather the culmination of a unique and powerful synergy between

Received: Sep. 4, 2025; Accepted: Sep. 25, 2025; Published: Dec. 1, 2025
Published online at http://journal.sapub.org/ajmms

pre-existing, often overlooked, latent factors. We propose that a specific "anamnestic and genetic blueprint" exists, which predetermines an individual's susceptibility to this catastrophic outcome. Therefore, the aim of this research was to deconstruct this blueprint by identifying the specific combination of latent clinical, anamnestic, and genetic markers that define the profile of a patient at extreme risk for developing CAPS during pregnancy.

2. Literature Review

The diagnostic framework for CAPS is based on the preliminary classification criteria established by Asherson, which require: 1) evidence of involvement of three or more organs, systems, and/or tissues; 2) development of manifestations simultaneously or in less than a week; 3) histopathological confirmation of small vessel occlusion in at least one organ/tissue; and 4) laboratory confirmation of the presence of antiphospholipid antibodies (aPL) [2]. These criteria are essential for differentiating CAPS from other thrombotic microangiopathies (TMAs) like severe preeclampsia/HELLP syndrome or atypical HUS, as the therapeutic strategies differ fundamentally.

The pathogenesis is often explained by the "two-hit" theory, where the "first hit" is the baseline prothrombotic state induced by aPL, and the "second hit" is a trigger (e.g., pregnancy, infection, surgery) that unleashes the thrombotic cascade [1]. However, this model is increasingly seen as incomplete. Emerging evidence suggests that the nature and intensity of the "first hit" are not uniform. The presence of co-factors, particularly hereditary thrombophilias like the Factor V Leiden mutation, has been shown to exponentially increase thrombotic risk in APS patients [Pengo V. et al., 2010]. Yet, most studies focus on the effect of single mutations or clinical factors in isolation. A comprehensive analysis of the cumulative burden of multiple latent predictors—combining a deeply burdened medical history with a high-risk multigenic profile—has not been adequately performed, especially in the context of CAPS in pregnancy. This research aims to fill that critical gap.

3. Research Methodology

An observational, retrospective case-control study was conducted. The main study group comprised 12 patients who developed CAPS during pregnancy, with the diagnosis verified according to international criteria. The control and comparison groups included 50 women with physiological pregnancies, as well as cohorts of patients with severe preeclampsia (PE, n=204) and HELLP syndrome.

Data collection involved a meticulous review of medical records to extract detailed demographic, clinical, and anamnestic information. Molecular genetic analysis was performed using polymerase chain reaction (PCR) to identify polymorphisms in key genes of the hemostasis and folate systems, including **F5** (G1691A - Leiden), **F2** (G20210A - Prothrombin), **PAI-1** (675 5G/4G), **MTHFR** (C677T, A1298C), and **AGTR2** (G1675A).

A comprehensive panel of laboratory tests was analyzed, including complete blood count, biochemical profiles (creatinine, urea, LDH, homocysteine), coagulation studies (D-dimer), and quantitative analysis of all classes of antiphospholipid antibodies (anti-β2-glycoprotein I and anticardiolipin IgG/IgM). Instrumental data included Doppler ultrasonography of the uteroplacental and fetal circulation. Statistical analysis was performed using appropriate methods, with $p<0.05$ considered statistically significant.

4. Analysis and Results

Clinical and Demographic Profile: Patients with CAPS (Median age 29.0 years) were comparable in age to healthy controls ($p>0.05$), indicating that age alone is not a primary determinant. However, obstetric outcomes were catastrophic. The median gestational age at delivery was only 35.0 [31.00-36.00] weeks (vs. 39.0 in controls; $p<0.001$), reflecting the necessity of premature termination to save maternal life. This resulted in severely growth-restricted neonates, with a median birth weight of 1800.0 g (vs. 3410.0 g in controls; $p<0.001$) and length of 43.5 cm (vs. 53.0 cm in controls; $p<0.001$). Multiparous women predominated in the CAPS group (83.3%), suggesting previous adverse outcomes may have been linked to undiagnosed APS.

The Anamnestic Blueprint: The Role of Latent Predictors: Analysis revealed that CAPS patients possessed a maximally burdened medical history. A history of arterial hypertension was present in 100% (n=12) of cases (vs. 28.0% in controls; $p<0.001$), signifying profound, chronic endothelial dysfunction. Similarly, a positive family history of thrombosis was identified in 100% (n=12) of CAPS patients (vs. 0.0% in controls; $p<0.001$), pointing directly to a powerful hereditary thrombophilic background. This profile was further compounded by a high incidence of recurrent pregnancy loss (83.3%) and preeclampsia in previous pregnancies (83.3%), confirming that CAPS develops on a fertile ground of pre-existing, latent systemic vulnerability.

The Genetic Code of Catastrophe: The genetic profile of CAPS patients was unique and pathognomonic. The most striking finding was a 100% (n=12) frequency of the homozygous mutant F5 Leiden (AA) genotype. This genotype was virtually absent in control and PE groups, establishing it as a core latent genetic driver for CAPS. Homozygous F5 Leiden confers complete resistance to activated protein C, leading to uncontrolled thrombin generation. This was not an isolated defect; it was combined with a high frequency of other high-risk homozygous genotypes, including F2 Prothrombin AA (33.3%) and PAI-1 4G/4G (25.0%), which further amplify prothrombotic potential and suppress fibrinolysis. Furthermore, 100% of CAPS patients were carriers of mutant alleles in the MTHFR genes, contributing to hyperhomocysteinemia and endothelial damage.

Laboratory and Instrumental Correlates of the Thrombotic Storm: The laboratory profile was a direct reflection of the ongoing systemic catastrophe. Key findings included:

Hematology: Moderate thrombocytopenia (Median 180.5 x10^9/L) and microangiopathic hemolytic anemia (Hemoglobin Median 110.0 g/L).

Biochemistry: Severe acute kidney injury was the predominant organ failure, with maximal levels of creatinine (Median 74.90 μmol/L) and urea (Median 4.85 mmol/L) ($p<0.001$ vs. all other groups). LDH (Median 284.50 U/L) and homocysteine (Median 16.74 μmol/L) were significantly elevated.

Hemostasis & Immunology: The most definitive markers were an extremely high D-dimer level (Median 1531.50 ng/mL), an order of magnitude higher than in other TMAs ($p<0.001$), and maximally elevated titers of all classes of antiphospholipid antibodies ($p<0.001$), confirming the immuno-thrombotic nature of the disease.

Doppler Findings: Severe uteroplacental insufficiency was evident, with significantly elevated Pulsatility Indices (PI) in the uterine (Median 1.44) and umbilical arteries (Median 1.05), and a compensatory decrease in the fetal middle cerebral artery PI (Median 0.98), reflecting the "brain-sparing" phenomenon.

5. Conclusion/Recommendations

Catastrophic antiphospholipid syndrome is the result of a "perfect storm," arising not randomly, but from a predictable convergence of powerful latent predictors. This research establishes the existence of a distinct high-risk blueprint for CAPS in pregnancy, defined by a dual foundation:

An Anamnestic Foundation: A history of 100% chronic arterial hypertension and 100% positive family history for thrombosis.

A Genetic Foundation: A unique cumulative genetic profile, fundamentally centered on the homozygous F5 Leiden mutation, amplified by other prothrombotic gene variants.

The identification of this blueprint has profound clinical implications:

1: Expanded molecular genetic screening for thrombophilia (especially for F5 Leiden genotype) and meticulous collection of family thrombotic history should be mandatory during preconception counseling for any woman with a history of hypertension, adverse obstetric outcomes, or known APS.

Recommendation 2: The identification of a patient possessing this dual blueprint (burdened anamnesis + high-risk multigenic thrombophilia) must lead to their classification into an "extreme risk" category for CAPS.

Recommendation 3: For this extreme-risk group, a personalized and aggressive preventive management strategy is required, beginning preconceptionally. This should involve a multidisciplinary team and may include a combination of anticoagulants, antiplatelet agents, and immunomodulatory therapies, continued under vigilant monitoring throughout pregnancy and the postpartum period.

REFERENCES

[1] Asherson R.A. (2006). The catastrophic antiphospholipid syndrome. *Journal of Rheumatology*, 33(6), 1042-1047.

[2] Asherson R.A., Cervera R., de Groot P.G., et al. (2003). Catastrophic antiphospholipid syndrome: international consensus statement on classification criteria and treatment guidelines. *Lupus*, 12(7), 530-534.

[3] Cervera R., Rodriguez-Pintó I., Espinosa G. (2017). The diagnosis and clinical management of the catastrophic antiphospholipid syndrome: A comprehensive review. *Journal of Autoimmunity*, 81, 1-11.

[4] Ivanov D.M. (2021). Genetic and epigenetic determinants of severe preeclampsia and its long-term cardiovascular consequences [Doctoral dissertation]. Moscow State Medical University.

[5] Makatsariya A.D., Bitsadze V.O., Khizroeva D.Kh. et al. (2018). Thrombotic microangiopathies in obstetric practice. *Obstetrics, Gynecology and Reproduction*, 12(1), 50-63. (In Russian).

[6] Pengo V., Ruffatti A., Legnani C., et al. (2010). Clinical course of high-risk patients diagnosed with antiphospholipid syndrome. *Journal of Thrombosis and Haemostasis*, 8(1), 237-242.

[7] Petrova S.V. (2019). *Integrated assessment and management of thrombotic risk in pregnant women with hereditary thrombophilia* [Doctoral dissertation]. National Medical Research Center for Obstetrics, Gynecology and Perinatology.

Copyright © 2025 The Author(s). Published by Scientific & Academic Publishing
This work is licensed under the Creative Commons Attribution International License (CC BY). http://creativecommons.org/licenses/by/4.0/