

Association Between Elevated Serum TNF- α Levels in Patients with Diabetes Mellitus and the Development of Diabetic Neuropathy

Saatov T. S.¹, Ibragimova E. A.¹, Toshtemirov A. E.¹, Alimov T. R.^{1,2}, Ibragimov Z. Z.¹,
Ishankhodjayev T. M.¹, Artykbaeva G. M.¹, Mustafakulov M. A.¹, Shamansurova Z. M.^{1,3},
Abdurakhimov S. A.¹, Nasriddinova P. M.¹, Maksutova N. N.⁴

¹Institute of Biophysics and Biochemistry at the National University of Uzbekistan named after Mirzo Ulugbek, Tashkent, Uzbekistan

²Republican Specialized Scientific-Practical Medical Center of Hematology, Ministry of Health of the Republic of Uzbekistan

³Central Asian University, Tashkent, Uzbekistan

⁴Republican Specialized Scientific-Practical Medical Center of Endocrinology named after academician Ya.Kh. Turakulov, Ministry of Health of the Republic of Uzbekistan

Abstract To investigate the relationship between TNF- α levels and the development of neuropathy in diabetes mellitus. The study involved patients with type 2 diabetes mellitus (T2DM) without complications and patients with diabetic neuropathy (DN) undergoing inpatient treatment at the Republican Specialized Scientific-Practical Medical Center of Endocrinology (Ministry of Health of the Republic of Uzbekistan). Biochemical parameters and TNF- α cytokine levels were measured in the blood. It was established that serum TNF- α levels in patients with diabetic neuropathy correlate directly with the severity of the disease.

Keywords T2DM, Diabetic neuropathy (DN), Cytokine, TNF- α

1. Introduction

In recent years, the role of the cytokine tumor necrosis factor- α (TNF- α) in the pathogenesis of various human diseases has been intensively studied. It has been established that TNF- α acts as a key pro-inflammatory mediator involved in the development of insulin resistance and the pathogenesis of type 2 diabetes mellitus. TNF- α is one of the triggering factors in the development of obesity, diabetes mellitus, and its complications [5,7]. A link has been identified between changes in TNF- α levels and the course of various metabolic, autoimmune, and neurodegenerative diseases [1,3].

According to the literature, elevated blood concentrations of TNF- α in diabetic patients may be associated with increased secretion of this cytokine by monocytes/macrophages in patients with type 2 diabetes [2].

TNF- α is detectable in the blood at various stages of the inflammatory process. Numerous studies indicate that elevated blood TNF- α concentrations serve as an indicator of carbohydrate metabolism disorders, impaired glucose tolerance, and insulin resistance in diabetes mellitus [10]. Evidence exists regarding the pro-atherogenic properties of

TNF- α and its role in lipid metabolism and the development of atherogenic dyslipidemia [8,9,15]. TNF- α has been shown to influence plasma triglyceride concentrations, contributing to decreased HDL-cholesterol levels and increased LDL-cholesterol levels. Elevated TNF- α levels block insulin signaling via serine phosphorylation of insulin receptor substrate-1 (IRS-1) and induce insulin resistance in adipocytes and peripheral tissues. It has been established that TNF- α contributes to the impairment of insulin's biological action, leading to the development of type 2 diabetes mellitus [4,13].

Diabetic neuropathy (DN) is a severe complication of diabetes mellitus. Assessing peripheral blood TNF- α levels may be significant in evaluating the progression of complications in type 2 diabetes, particularly diabetic neuropathy.

2. The Aim of the Study

The aim of the study was to investigate the relationship between serum TNF- α levels and the development of neuropathy in patients with diabetes mellitus (DM).

3. Materials and Methods

The study involved 70 patients with type 2 diabetes

mellitus undergoing inpatient treatment at the Academician Ya.Kh. Turakulov Republican Specialized Scientific-Practical Medical Center of Endocrinology (Ministry of Health of the Republic of Uzbekistan). Patients were divided into groups based on clinical history and clinical-laboratory findings:

Group I – patients with DM without complications (n=20)

Group II – patients with diabetic neuropathy (DN) (n=50)

Group III – individuals without DM (n=10)

Group II was further divided into two subgroups based on DN severity: moderate (DN1) and severe (DN2) disease course. The control group consisted of individuals with no clinical or laboratory signs of carbohydrate metabolism disorders.

Fasting capillary blood glucose levels were determined using the glucose oxidase method with "Cypress Diagnostics" (Belgium) kits. Biochemical analyses were performed using HUMAN (Germany) kits. Serum TNF- α concentrations in patients with DM were measured via enzyme-linked immunosorbent assay (ELISA) using "TNF-alpha-ELISA BEST" (Russia) kits.

Measurements were performed using an MR96 microplate photometer (Mindray, China).

4. Results and Discussion

Assessment of the subjects' age revealed no statistically significant difference between the group of patients with diabetes mellitus and the control group ($p > 0.05$). The duration of the disease was 7.67 ± 5.4 years.

Analysis of peripheral blood biochemical parameters in patients with type 2 diabetes mellitus (T2DM) yielded the following findings. Patients diagnosed with T2DM exhibited carbohydrate and lipid profiles characteristic of this pathology. A significant increase was observed in fasting glucose levels (3.6-fold) and glycated hemoglobin (HbA1c) levels (3.1-fold) ($p < 0.05$) compared to the control group. Regarding lipid abnormalities, the most significant changes

included a 3.1-fold increase in triglyceride (TG) levels ($p < 0.05$) and elevated very-low-density lipoprotein cholesterol (VLDL-C) levels; the latter may be attributed to either increased production or reduced catabolism of VLDL-C.

It is possible that the alterations in blood carbohydrate and lipid levels observed in T2DM patients were also associated with elevated levels of the cytokine TNF- α . Considering this possibility, the TNF- α levels in the peripheral blood of the study groups were measured.

ELISA-based analysis of serum TNF- α levels revealed a significant increase in TNF- α concentration in patients with diabetes (3.05 ± 0.44 pg/mL) compared to non-diabetic subjects (1.10 ± 0.05 pg/mL; $p < 0.05$).

A comparison of serum TNF- α levels in patients with type 2 diabetes mellitus—with and without diabetic neuropathy—revealed a trend toward increased TNF- α concentrations in patients who had developed diabetic neuropathy; the mean level across the entire patient group was 5.2 ± 0.44 pg/mL (Figure 1). Our study results showed no significant difference in TNF- α levels between patients with moderate diabetic neuropathy (DN1) and those without neuropathy (3.62 ± 0.89 pg/mL and 3.05 ± 0.44 pg/mL, respectively).

According to the results obtained, serum TNF- α levels in patients with diabetic neuropathy increased with the progression of the complication. Specifically, in patients with severe diabetic neuropathy, the TNF- α level was nearly twice as high as in those with moderate diabetic neuropathy, measuring 7.13 ± 1.14 pg/mL and 3.62 ± 0.89 pg/mL, respectively (Figure 2).

Thus, the results indicate a significant increase in serum TNF- α concentration in patients with type 2 diabetes mellitus (T2DM). A statistically significant rise in TNF- α levels is observed during the development of complications, particularly in cases of severe diabetic neuropathy. It has been established that the serum concentration of this cytokine in patients with diabetic neuropathy correlates directly with the severity of the disease.




Figure 1. Comparison of serum TNF- α levels in patients with type 2 diabetes mellitus without neuropathy and with diabetic neuropathy




Figure 2. Serum TNF- α levels in patients with diabetic neuropathy (pg/mL, $p < 0.05$); DN – combined group of patients with diabetic neuropathy; DN1 – moderate diabetic neuropathy; DN2 – severe diabetic neuropathy

Our findings are supported by several studies [6,11,12]. Elevated serum TNF- α levels in patients with diabetes may be linked to the activation of inflammatory processes, the development of complications, and the worsening of disease severity. Specifically, higher levels of this cytokine were found in patients with complicated forms of diabetes compared to those with uncomplicated forms. The rise in serum TNF- α in T2DM patients is accompanied by lipid metabolism disorders, consistent with data regarding the pro-atherogenic properties of this cytokine and its role in the development of atherogenic dyslipidemia [14].

To date, no biochemical markers have been established for the early diagnosis and monitoring of neuropathy progression. In our view, TNF- α could serve as an effective biochemical marker for the early diagnosis of diabetic neuropathy.

In recent years, research has actively focused on identifying inhibitors to block the adverse effects of TNF- α [4]. The development of such specific inhibitors holds promise for the treatment of insulin resistance and type 2 diabetes.

5. Conclusions

It has been established that serum TNF- α levels in patients with diabetes mellitus correlate directly with the severity of diabetic neuropathy. Measuring peripheral blood TNF- α levels may have prognostic value in assessing the progression of type 2 diabetes complications, particularly diabetic neuropathy.

REFERENCES

- [1] Alesenko A.V., Bachurin S.O., Guryanova S.V., Karatasso Yu.O., Shevtsova E.F., Shingarova L.N. Tumor necrosis factor-alpha: a potential target for the neuroprotective drug Dimebon // *Biomedical Chemistry*. 2016, Vol. 62, Issue 4, pp. 418–425.
- [2] Voronina E.V., Lobanova N.V., Yakhin I.R., Romanova N.A., Seregin Yu.A. The role of tumor necrosis factor-alpha in the immunogenesis of diseases of various etiologies and its significance in the development of anti-TNF monoclonal antibody therapy // *Medical Immunology*. 2018, Vol. 20, No. 6, pp. 797–806.
- [3] Mirzoeva L.A., Nikiforov N.G., Aladinsky V.A., Sobenin I.A., Nedosugova L.V., Orekhov A.N. Increased spontaneous and induced secretion of the pro-inflammatory cytokine TNF- α by blood monocyte-macrophages in patients with type 2 diabetes mellitus. *Problems of Endocrinology*. – 2014. – Vol. 60. – No. 5. – P. 22–25.
- [4] Akash M.S.H., Rehman K., Liaqat A. Tumor Necrosis Factor-Alpha: Role in Development of Insulin Resistance and Pathogenesis of Type 2 Diabetes Mellitus. // *J. Cell Biochem*. 2018 Jan; 119(1): 105–110.
- [5] Bashir, H., Ahmad Bhat, S., Majid, S., Hamid, R., Koul, R. K., Rehman, M. U., Din, I., Ahmad Bhat, J., Qadir, J., & Masood, A. (2020). Role of inflammatory mediators (TNF- α , IL-6, CRP), biochemical and hematological parameters in type 2 diabetes mellitus patients of Kashmir, India. *Medical journal of the Islamic Republic of Iran*, 34, 5.
- [6] Chen, L., Chen, R., Wang, H., & Liang, F. (2015). Mechanisms Linking Inflammation to Insulin Resistance. *International journal of endocrinology*, 2015, 508409.
- [7] Chenxiao Liu C., Feng X., Li Q., Wang Y Li Q., Hua M. Adiponectin, TNF-a and inflammatory cytokines and risk of type 2 diabetes: A systematic review and meta-analysis // *Cytokine* 86 (2016) 100–109.
- [8] Cure O, Kizilkaya B, Ciftel S, Mercantepe F. The Effect of Anti-Tumor Necrosis Factor Therapy on The Plasma Atherogenic Index in Rheumatic Diseases. *Metabolic Syndrome and Related Disorders*. 2024; 22(4): 269–275.
- [9] Fischer, R., & Maier, O. (2015). Interrelation of oxidative stress and inflammation in neurodegenerative disease: role of

- TNF. Oxidative medicine and cellular longevity, 2015, 610813.
- [10] Moriwaki Y, Yamamoto T, Shibutani Y, Aoki E, Tsutsumi Z, Takahashi S, Okamura H, Koga M, Fukuchi M, Hada T. Elevated levels of interleukin-18 and tumor necrosis factor- α in serum of patients with type 2 diabetes mellitus: relationship with diabetic nephropathy. *Metabolism*. 52 (2003) 605-608.
- [11] Navarro J.F., Carmen Mora, Marina Gormez, Mercedes Muros, Celeste Lopez-Aguilar and Javier García Influence of renal involvement on peripheral blood mononuclear cell expression behaviour of tumour necrosis factor- α and interleukin-6 in type 2 diabetic patients. *Nephrol Dial Transplant* (2008) 23: 919-926.
- [12] Shamansurova Z.M., Saatov T.S., Takhrirov L.Sh. Tumor necrosis factor alpha: Role in the development of obesity and diabetes mellitus. *Asian Journal of Biocemistry, Genetics and Molecular Biology*. 2020. Jul. 4(3) 29-42.
- [13] Sookoian SC, González C, Pirola CJ. Meta-analysis on the G-308A tumor necrosis factor alpha gene variant and phenotypes associated with the metabolic syndrome. *Obes Res*. 2005; 13(12): 2122-2131.
- [14] Yang, Y., Zhao, B., Wang, Y., Lan, H., Liu, X., Hu, Y., & Cao, P. (2025). Diabetic neuropathy: cutting-edge research and future directions. *Signal transduction and targeted therapy*, 10(1), 132. <https://doi.org/10.1038/s41392-025-02175-1>.
- [15] Kubaszek A, Pihlajamäki J, Komarovski V, et al. Promoter polymorphisms of the TNF- α (G-308A) and IL-6 (C-174G) genes predict the conversion from impaired glucose tolerance to type 2 diabetes: the Finnish Diabetes Prevention Study. *Diabetes* 2003 Jul; 52(7): 1872-1876.