

Changes in Regional Lymph Node Morphology in Secondary Acute Peritonitis

E. A. Eshbaev^{1,*}, A. Q. Xaydarov²

¹Tashkent State Medical University, Tashkent, Uzbekistan

²Ferghana Medical Institute of Public Health, Fergana, Uzbekistan

Abstract Acute secondary peritonitis is the main source of morphological changes in the mesenteric lymph nodes in 18-45 year olds, which comes with relaxation of the morphofunctional areas of the lymph nodes in the first 1-3 days of peritonitis. It is determined that the development of lymph node lesions in the cortex and medulla is accompanied by the presence of lymphadenitis, various fragmented cell fragments, coarse colloidal fluid, macrophages, a small number of neutrophils, and the appearance of lymphadenitis. At the same time, the morphological features are characterized by the presence of large macrophages undergoing incomplete phagocytosis in the reticular fibers of the subcapsular and medulla layers of the lymph nodes. It is precisely these macrophages that are found in the form of adhesions to the interdigitating cells of the medulla layer of the lymph node.

Keywords Lymph node, Peritonitis, Lymphoid follicle, Immunodeficiency

1. Introduction

In the world today, the lethality of purulent peritonitis is very high, reaching 20-60% [1,2,3]. When this disease occurs, it is accompanied by the development of rapidly developing endogenous intoxication and multi-organ failure. In the USA and Europe, this indicator is 10-15 per 100 thousand, in Turkey and Korea it is 12-17 per 100 thousand. In the Russian Federation and the CIS countries it is 15-20 per 100 thousand, in the countries of Central Asia, Azerbaijan, Armenia this indicator is 50-71 per 100, with a lethality of 60-85% [4,5,6,7,8]. In our country, on average, 700-1300 infants under the age of 1 die from various types of peritonitis per year. In peritonitis [9,10], the main organs actively involved in the drainage of the abdominal cavity are only the regional lymph nodes, and at 1-3 months of age, this drainage is dominated by the occurrence of physiological paralysis and slow blood circulation, while at 3-6 months, the weakening of intestinal peristalsis and drainage in the regional lymph nodes, insufficiency of the barrier function, lead to the stagnation of local infection in the intestinal lymph nodes and the development of lymphostasis. This requires the development of specific recommendations, taking into account the specific morphofunctional aspects of each period [11,12,13,14].

2. Materials and Methods

A total of 131 patients were recruited to the Republican Center for Pathological Anatomy in 2018-2023, including 15 patients who died of acute myocardial infarction as a control group. The obtained samples were stained with hematoxylin and eosin and studied.

3. Objective

To study and improve the morphological features of the development of mesenteric lymph nodes in acute secondary peritonitis in people aged 18-45.

4. Results and Discussion

Acute secondary peritonitis of various etiologies, the response of regional lymph nodes during the first 24-72 hours proceeds according to the following morphological changes.

In acute purulent peritonitis, as a result of the drainage of inflammatory exudate through the peritoneal surfaces through massive lymphatic vessels, stasis and congestion of lymphatic vessels and edema of lymph nodes occur. The main accumulated product is fragments consisting of inflammatory exudate and disintegrated cellular components. Thus, a large amount of pathological fluid is accumulated in the subcapsular areas of the lymph node, a large amount of tissue macrophages are accumulated in the lymphatic vessels and subcapsular areas of the lymph node, the blood vessels of the portal vein of the lymph node are filled, massive

* Corresponding author:

eshbaev@yahoo.com (E. A. Eshbaev)

Received: Sep. 12; Accepted: Oct. 8, 2026; Published: Oct. 30, 2026

Published online at <http://journal.sapub.org/ijvmb>

migration of immune cells to the foci in the cortical and paracortical areas, in particular, the mobilization of antigen-dependent differentiated lymphocytes from the secondary follicles formed in the germinal center of lymphoid follicles through postcapillary venules, and the **germinal centers are exposed and empty**.




Figure 1. A small image of the mesenteric lymph node shows signs of massive perfusion in all blood vessels. A massive decrease in lymphocytes in the stroma, foci of delymphatization in the cortical layer. Coarsely dispersed pink homogeneous deposits are detected in the subcapsular area. Staining GE. Size 4x10

These changes are characterized by a sharp decrease in T-lymphocytes in the paracortical areas and an increase in reticulocytes, macrophages and a small amount of fibroblasts. This means that in acute peritonitis, within the first 24 hours, all antigen-specific differentiated lymphocytes in the regional lymph nodes migrate to the foci of inflammation. At the same time, in the medullary layer of the lymph node, a variable mixture of T and B lymphocytes in the medullary bands, the ratio of histiocytes and reticulocytes changes sharply. Mainly, an accumulation of reticulocytes, fibroblasts, histiocytes and macrophages is detected. Intermediate tumors also develop in the medullary layer, which indicates the accumulation of various dispersed metabolites in large quantities, and the development of acute morphofunctional disorders in the lymph nodes. This, in turn, leads to the following changes in the morphofunctionally active areas of the lymph node: The lymph nodes appear swollen, their capsule is tense, and the subcapsular spaces are filled with lymphatic fluid.




Figure 2. Mesenteric lymph node. Macrophages and reticulocytes are abundantly accumulated and lymphocytes are sharply reduced. Paint G.E. The size is 40x10

Secondary lymphoid follicles are not detected, the germinal center is stable, the active areas of the lymph nodes are less densely populated than usual, the localization of lymphocytes in the cortical and paracortical areas of the active areas of the cortex is reduced, and in the active areas of the cortex, foci of lymphocytes migrating towards the postcapillary venules are detected. In the direction of the medullary cords, the lymphatic portal areas are edematous, and in the reticular area and trabeculae of the medullary cords, a variable mixture of lymphocytes and other types of cells (macrophages, dendritic cells, interdigitating, fibroblasts, histiocytes, reticulocytes, etc.) is relatively preserved. Within 1 day of peritonitis development, the peritoneum shows predominantly venous congestion, lymphostasis, multifocal desquamation of mesothelial cells, and foci of mucoid thickening of the basement membrane of the exposed sheets. The active areas of the lymph nodes are blurred, and foci of sharply migrating lymphocytes are detected in the postcapillary venules. The lymphatic vessels are full and filled with macrophages.




Figure 3. Mesenteric lymph node. In the subcapsular space, macrophages and various tissue fragments, cell fragments are detected (1), pink homogeneous protein deposits and Mott cells are detected (2). Paint G.E. 10x10

It is precisely because the main part of the regional lymph nodes is located in the ileocecal mandible and precisely in peritonitis, massive enlargement of the lymph nodes proceeds. The capsules of the lymph nodes are taut, uniform thickness, also, reticulocytosis and macrophages can be detected in the subcapsular space.




Figure 4. Coarse homogeneous protein deposits, tissue fragments are detected in subcapsular spaces. Macrophages and Mott cells are detected (1). Paint G.E. The size is 40x10

As a result of a sharp increase in the level of fluid drainage, massively exposed stroma is detected in the lymphoid follicles in the active areas of the lymph node (cortical and paracortical areas).

Lymphoid follicles are sharply blurred against the general background, the follicle texture is lost, and foci of congestion and stasis are detected in the postcapillary venules in the paracortical area.

In the medullary layer, a variable admixture of other cell types, predominantly macrophages, and an increase in mast cells are detected in the direction of the cerebral striations. These indicators determine the dynamics of the inflammatory process. These changes are accompanied by the development of acute inflammation, vasoparalysis and vasodilation (acute expansion) of the vessels in response to inflammatory mediators, increased absorption of metabolites through paralysis of the vascular wall, clinically, toxic substances cause toxic dystrophy in parenchymal organs (liver, kidneys, heart, lungs) and ultimately lead to the process of endogenous intoxication and multiorgan failure. It is important to note that this is characterized by the rapid degranulation of mast cells and neutrophils and the release of large amounts of cytokines, leading to a "cytokine storm," which leads to a systemic vascular response and ultimately to multiorgan failure.




Figure 5. Mesenteric lymph node. Macrophages and reticulocytes are abundantly accumulated and lymphocytes are sharply reduced. Paint G.E. Size 4x10

Clinically, it is characterized by severe swelling of the organs, diapedesis, hemorrhages, the appearance of fibrous plaques, foci of microabscesses on the serous membrane surfaces.

At the same time, there is an increase in mast cells, an admixture of destroyed fibrous structures and tissue detritus, and histologically, dispersed debris formations are preserved. Lymphostasis in the lymphatic vessels, dilation of lymphatic capillaries are preserved.

5. Conclusions

Acute secondary peritonitis is the main source of morphological changes in the mesenteric lymph nodes in individuals aged 18–45 years, manifested by relaxation of the morphofunctional zones of the lymph nodes during the first 1–3 days of the disease. The development of lesions in the cortex and medulla is accompanied by the presence of

lymphadenitis, various fragmented cell fragments, coarsely dispersed colloidal fluid, macrophages, a small number of neutrophils, and the appearance of lymphadenitis. At the same time, the morphological features are characterized by the presence of large macrophages undergoing incomplete phagocytosis in the reticular fibers of the subcapsular and medullary layers of the lymph nodes. It is precisely these macrophages that are found in the form of adhesions to the interdigitating cells of the medullary layer of the lymph node.

Within the first 24 hours of peritonitis development, all antigen-specific differentiated lymphocytes in the regional lymph nodes migrate to the foci of inflammation. This is accompanied by a sharp decrease in T-lymphocytes in the paracortical areas and an increase in reticulocytes, macrophages, and a small amount of fibroblasts. In the medullary layer, a variable mixture of T and B lymphocytes in the medullary bands is observed, and the ratio of histiocytes and reticulocytes changes sharply. Mainly, an accumulation of reticulocytes, fibroblasts, histiocytes, and macrophages is detected. Intermediate tumors also develop in the medullary layer, which indicates the accumulation of various dispersed metabolites in large quantities and the development of acute morphofunctional disorders in the lymph nodes. The lymph nodes appear swollen, their capsule is tense, and the subcapsular spaces are filled with lymphatic fluid. Secondary lymphoid follicles are not detected, the germinal center is stable, and the active areas of the lymph nodes are less densely populated than usual. Foci of lymphocytes migrating toward the postcapillary venules are detected in the active areas of the cortex, and the lymphatic portal areas are edematous in the direction of the medullary cords.

As a result of a sharp increase in the level of fluid drainage, massively exposed stroma is detected in the lymphoid follicles in the active areas of the lymph node. Lymphoid follicles are sharply blurred against the general background, the follicle texture is lost, and foci of congestion and stasis are detected in the postcapillary venules in the paracortical area. In the medullary layer, a variable admixture of other cell types, predominantly macrophages, and an increase in mast cells are detected in the direction of the cerebral striations. These changes are accompanied by the development of acute inflammation, vasoparalysis, and vasodilation of the vessels in response to inflammatory mediators, increased absorption of metabolites through paralysis of the vascular wall. Clinically, toxic substances cause toxic dystrophy in parenchymal organs and ultimately lead to endogenous intoxication and multiorgan failure. This is characterized by the rapid degranulation of mast cells and neutrophils and the release of large amounts of cytokines, leading to a "cytokine storm," which leads to a systemic vascular response and ultimately to multiorgan failure. Clinically, it is characterized by severe swelling of the organs, diapedesis, hemorrhages, the appearance of fibrous plaques, and foci of microabscesses on the serous membrane surfaces. At the same time, there is an increase in mast cells, an admixture of destroyed fibrous

structures and tissue detritus, and histologically, dispersed debris formations are preserved. Lymphostasis in the lymphatic vessels and dilation of lymphatic capillaries are preserved. Thus, the morphological features of regional lymph node lesions in secondary acute peritonitis reflect the severity of the inflammatory process and play an important role in the development of endogenous intoxication and multiorgan failure, which requires the development of specific recommendations taking into account the specific morphofunctional aspects of each period of the disease.

REFERENCES

- [1] Gorbatyuk O. M., Martynyuk T. V., Shatrova K. M. Clinical and morphological characteristics of gastrointestinal perforations in newborns // *Russian Bulletin of Pediatric Surgery, Anesthesiology and Reanimatology*. – 2013. – Vol. 3. – No. 2. – P. 31-36.
- [2] Gorbatyuk O. M., Shatrova K. M., Martynyuk T. V. Morphological diagnosis of gastrointestinal perforations in newborn infants // *Neonatology, Surgery and Perinatal Medicine*. – 2015. No. 5, No. 3. P. 62-66.
- [3] Aires J, Ilhan ZE, Nicolas L, Ferraris L, Delannoy J, Bredel M, Chauvire-Drouard A, Barbut F, Roz éJC, Lepage P, Butel MJ, ClosNEC Study Group. Occurrence of Neonatal Necrotizing Enterocolitis in Premature Neonates and Gut Microbiota: A Case-Control Prospective Multicenter Study. // *Microorganisms*. 2023 Sep 29; 11(10): 2457.
- [4] Alsaied A, Islam N, Thalib L. Global incidence of necrotizing enterocolitis: a systematic review and meta-analysis. // *BMC Pediatr* 2020; 20: 344.
- [5] Andrade JAB, Freymüller E, Fagundes-Neto U. Pathophysiology of enteroaggregative *Escherichia coli* infection: an experimental model utilizing transmission electron microscopy. // *Arq Gastroenterol*. 2010; 47: 306–12.
- [6] Aujla S.J., Chan Y.R., Zheng M., Fei M., Askew D.J., Pociask D.A., Reinhart T.A., McAllister F., Edeal J., Gaus K. IL-22 mediates mucosal host defense against Gram-negative bacterial pneumonia. *Nat. Med.* 2008; 14: 275–281.
- [7] Aydemir G, Cekmez F, Tanju IA, Canpolat FE, Genc FA, Yildirim S, Tunc T, Sarici SU. Increased fecal calprotectin in preterm infants with necrotizing enterocolitis. // *Clin Lab*. 2012; 58(7-8): 841-4.
- [8] Berry MJ, Port LJ, Gately C, Stringer MD. Outcomes of infants born at 23 and 24 weeks' gestation with gut perforation. // *J Pediatr Surg*. 2019 Oct; 54(10): 2092-2098.
- [9] Bethell GS, Knight M, Hall NJ. BAPS-CASS B-CNIGobo. Surgical necrotizing enterocolitis: association between surgical indication, timing, and outcomes. // *J Pediatr Surg*. 2021; 56(10): 1785–90.
- [10] Bhatia AM, Stoll BJ, Cismowski MJ, Hamrick SE. Cytokine levels in the preterm infant with neonatal intestinal injury. // *Am J Perinatol*. 2014; 31(6): 489–96.
- [11] Black RE, Cousens S, Johnson HL, et al. Global, regional, and national causes of child mortality in 2008: a systematic analysis. // *Lancet*. 2010; 375: 1969–87.
- [12] Brusselaers N, Simin J, E Lilja H. Risk of neurodevelopmental impairment in Swedish preterm children treated for necrotizing enterocolitis: retrospective cohort study. // *BJS Open*. 2024 Oct 29; 8(6): zrae131.
- [13] Brusselaers N, Simin J, E Lilja H. Risk of neurodevelopmental impairment in Swedish preterm children treated for necrotizing enterocolitis: retrospective cohort study. // *BJS Open*. 2024 Oct 29; 8(6): zrae131.
- [14] Burnand KM, Zapparackaite I, Lahiri RP, Parsons G, Farrugia MK, Clarke SA, DeCaluwe D, Haddad M, Choudhry MS. The value of contrast studies in the evaluation of bowel strictures after necrotising enterocolitis. // *Pediatr Surg Int*. 2016 May; 32(5): 465-70.