

Human Papillomavirus (HPV) in Oral Cavity Lesions: Comparison with Other Oral Cancer Risk Factors

Raúl Fernando Venezuela^{1,2,*}, Ángel Talavera³, María Celia Frutos¹, Ana Ximena Kiguen¹,
Marina Soledad Monetti¹, María Sollazo³, René Panico³, Ruth Ferreyra de Prato³,
Cecilia Gabriela Cuffini¹

¹Institute of Virology, National University of Córdoba. Córdoba, 5016, Argentina

²Enfermera Gordillo Gómez s/n Ciudad Universitaria, Córdoba, 5016, Argentina

³Faculty of Dentistry - National University of Córdoba, 5016. Argentina

Abstract Human papillomavirus (HPV) is considered a necessary factor for the development of cervical cancer; however, its relationship with oral cancer is controversial. The aim of this study was detect the presence of HPV in lesions of the oral cavity and its correlation with other risk factors. Presence of HPV was studied by polymerase chain reaction in samples from benign lesions, potentially malignant lesions (PML), neoplasias and healthy mucosae. The results from the different groups were compared; in addition to their histopathological variables with tobacco smoking, alcohol consumption, so on. HPV was detected in 88.89% of benign lesions, 41.38% of PML and 56.25% of neoplasias. The most prevalent genotypes were 16 and 6. Together, reached 55% of the total number of cases. A significant association was observed between HPV and male gender, tobacco smokers, alcohol drinkers and benign lesions. Tobacco smoking and alcohol intake were associated to neoplasias. Our results showed that factors like tobacco smoking and alcohol drinking, have more influence than HPV in the development of oral neoplasias; however, 56.2% of the neoplasias tested positive for HPV; the percentage of HR-HPV detection increased with the severity of the lesions, suggesting its possible involvement in malignant processes

Keywords HPV, Oral cavity lesions, Argentina

1. Introduction

Head and neck cancer is a major health problem worldwide that usually appears in patients older than 50 years of age; however some studies have shown that between 1 to 6 percent of oral malignant tumors have been found in patients younger than 40[1,2,3].

Head and neck cancer defines a heterogeneous group of malignant lesions that involve different sites with similar risk factors and pathological features. The majority of these malignancies are oral squamous cell carcinomas (OSCC) and are characterized by a multifactorial etiopathogenesis [4,5].

The causal association between tobacco consumption and alcohol intake with the development of OSCC is well established; however, a considerable proportion of OSCC occurs in non smokers and non drinkers, indicating the presence of other risk factors[5,6,7].

During the last decades, epidemiological and molecular data have indicated the involvement of high-risk human

papillomavirus (HR-HPV) in these diseases, which was first proposed by Syrjanen in 1983 and afterwards supported by other authors[5,8,9,10] on the basis of the epitheliotropic nature of HPV[11,12], in addition to the the widely confirmed oncogenic potential of HR-HPV in the pathogenesis of anogenital cancer, especially cervical squamous cell carcinoma[13,14] and the morphological similarities between oropharyngeal and genital epithelia [15].

Although HPV has been found in OSCC, the wide range of viral prevalence reported in the literature has not contributed to the clarification of the relationship between HPV and oral carcinogenesis[16,17].

The aim of this study was to determine the frequency of HPV detection and circulating genotypes in lesions of the oral cavity as well as their correlation to other risk factors.

2. Material and Methods

Patients: This case-control epidemiological study was performed in patients diagnosed and treated at the Faculty of Dentistry - National University of Córdoba (patients were included from March 2012 to December 2012 and control cases from October 2012 to December, 2012). All the

* Corresponding author:

venezuelafermando@gmail.com (Raúl Fernando Venezuela)

Published online at <http://journal.sapub.org/microbiology>

Copyright © 2013 Scientific & Academic Publishing. All Rights Reserved

patients underwent a careful analysis of their clinical records and signed a written informed consent before being included into the study.

The patients presented histopathological diagnosis of benign lesions (vegetative lesions, wart and condylomateous) potentially malignant lesions (leukoplakia, lichen, verrucous leukoplakia, keratoacanthoma)[18] and neoplasias (verruco us carcinoma, carcinoma in situ-CIS and OSCC). Control cases were individuals with healthy oral mucosae who attended the Faculty of Dentistry for tooth extractions (Control group).

The data were classified according gender, age (greater or younger than 50 years), conventional risk factors (smoking and drinking) site, histopathological diagnosis and cytological classification of the lesions.

Consideration of an individual as a smoker and/or drinker was based on the study of Herrero 2003[19].

The study population comprised 84 patients (9 with benign lesions, 29 potentially malignant lesions, 16 neoplasias and 30 control subjects).

Sample collection: All participants were subjected to an oral examination for the collection of cells (brushing) for HPV-DNA detection. Patients with presence of lesion, post-brushing for HPV detection and cytology diagnosis (Papanicolaou or PAP stains), underwent a biopsy which was collected in paraffin blocks. For control group was done brushing of full mouth.

The brushing samples for PCR were collected in 500 μ L phosphate-buffered saline solution (PBS).

HPV detection: Viral DNA was extracted using the commercial AccuPrep Genomic DNA Extraction Kit (Bioneer Inc., CA, USA), in accordance with the manufacturer's instructions.

A 450-bp segment, corresponding to the L1 region of the viral genome, was amplified by PCR, using the degenerate consensus primers MY09 and MY11 (Integrated DNA Technology - USA)[20]. The product was detected by electrophoresis in 1.5% agarose gel using a U.V. transilluminator. The β -globin gene was used as a DNA preservation marker[21]. Negative samples were considered inadequate.

HPV-DNA positive samples were typed by Restriction Fragment Length Polymorphism (RFLP) according to Bernard et al., 1994. Briefly, aliquots of the PCR products obtained using the degenerate consensus primers MY09–MY11, targeting a region of approximately 450 bp in length in the L1 ORF of the viral genome, were mixed with 7 different restriction enzymes (Bam HI, Hae III, Dde I, Pst I, Hinf I, Sau III, and Rsa I) in separate reactions. The digestion products were separated by electrophoresis in a 3% agarose gel and the pattern obtained was compared with published data.

Statistical analysis: we used the software InfoStat version 2011, with a significance level of 5% (95% CI) (InfoStat, computer program 2011)[22].

Chi-square (χ^2) tests were conducted to determine the

possible associations between presence, type and location of the lesions, cytological classification, history of tobacco smoking habit, alcohol consumption, age, gender, HPV and other risk factors.

3. Results

The patients were classified into four groups, as follows:

Control group: This group comprised 30 individuals, 8 males and 22 females, mean age: 41.8 ± 17.1 years (range, 18-81). Most controls were non tobacco smokers and non drinkers ((23/30 and 25/30, respectively). Inclusion criteria were: immunocompetent status and absence of any detectable lesions in the oral mucosa.

Patients with benign lesions: The mean age of the 9 patients of this group (1 female, 8 males) was 34.0 ± 16.8 years (range, 18-58 years); 2 patients were tobacco smokers, 2 reported regular alcohol consumption and 1 both habits. The most frequent locations were the buccal mucosa (n=3) and tongue (n=3), followed by uvula (n=1), hard palate (n=1) and gum. The distribution by cytology showed: grade II (n=7) and grade III (n=2).

Patients with potentially malignant lesions (PML): The mean age of the 29 patients (16 females, 13 males) was 52.6 ± 14.7 years (range, 29-76 years). Four patients were tobacco smokers, 4 reported regular alcohol consumption and 3 both habits. The most frequent location was tongue (n=18), followed by oral mucosa (n=8), hard palate (n=1), mouth floor (n=1) and lip (n=1). The results of cytology showed: grade II (n=10) and grade III (n=19).

Patients with neoplasia: The mean age of the 16 patients (8 females, 8 males) was 59.6 ± 19.7 years (range, 23-89 years), 10 were smokers; 7 reported alcohol consumption and 5 both habits. The most frequent location was tongue (n=8), followed by the oral mucosa (n=3), retromolar trigone (n=1), lip (n=1), gum (n=1), hard palate (n=1) and mouth floor (n=1). The distribution based on the cytological findings was: grade III (n=7), grade IV (n=3) and grade V (n=6).

HPV was detected in 88.89% (8/9) of the samples from oral benign lesions, 41.38% (12/29) of oral PML samples and 56.25% (9/16) of oral neoplasias. In samples from the control group, HPV was not detected.

Table 1 shows the characteristics of HPV-positive patients and the genotypes detected. The most prevalent were genotypes 16 and 6. Together, these two genotypes reached 55% of the total number of patients and both were identified within the 3 groups of patients. Figure 1 shows the percentage of detection of HR-HPV and LR-HPV in the different groups.

Table 2 shows the correlation between HPV and the different variables. A significant association of HPV with tobacco smoking habit, alcohol drinking and male gender was detected, but no association was found in patients over 50 years of age. With these data, we performed a statistical analysis between the different variables including the finding of HPV, versus clinical and histopathological diagnosis.




Figure 1. Detection of HR-HPV and LR-HPV in the different lesions

Table 1. Characteristics of HPV-positive patients and the genotypes detected

Age	Gender	Lesion	Histopathology	Trauma	Tobacco smoker	Alcohol drinker	Localization of Lesion	Genotype
24	M	benign	vegetative lesions	No	No	No	gum	6 LR-HPV
56	M	benign	wart	Yes	Yes	No	tongue	6 LR-HPV
18	M	benign	vegetative lesions	Yes	No	No	buccal mucous	6 LR-HPV
58	M	benign	wart	No	No	No	hard palate	6 LR-HPV
18	M	benign	condylomateous	No	No	Yes	tongue	13 LR-HPV
29	F	benign	vegetative lesions	No	Yes	Yes	buccal mucous	16 HR-HPV
29	M	benign	vegetative lesions	No	No	No	tongue	31 HR-HPV
53	M	benign	wart	No	No	No	buccal mucous	72 LR-HPV
65	M	PML	verrucous leukoplakia	Yes	No	Yes	tongue	6 LR-HPV
43	F	PML	verrucous leukoplakia	No	No	No	tongue	6 LR-HPV
47	M	PML	verrucous leukoplakia	No	No	No	buccal mucous	6 LR-HPV
47	F	PML	leukoplakia	Yes	No	No	buccal mucous	11 LR-HPV
63	F	PML	keratotic lichen	No	No	No	tongue	11 LR-HPV
41	M	PML	keratotic lichen	Yes	Yes	No	buccal mucous	13 LR-HPV
45	M	PML	verrucous leukoplakia	No	No	No	buccal mucous	16 HR-HPV
48	M	PML	keratoacanthoma	Yes	Yes	Yes	lip	16 HR-HPV
33	F	PML	leukoplakia	No	No	No	tongue	16 HR-HPV
55	F	PML	keratotic lichen	Yes	No	No	tongue	52 HR-HPV
35	F	PML	verrucous leukoplakia	No	No	No	buccal mucous	52 HR-HPV
53	M	PML	leukoplakia	No	Yes	Yes	buccal mucous	61 LR-HPV
77	M	neoplasia	oral squamous cell carcinoma	No	Yes	Yes	tongue	6 LR-HPV
63	M	neoplasia	oral squamous cell carcinoma	No	Yes	Yes	floor of mouth	11 LR-HPV
79	M	neoplasia	oral squamous cell carcinoma	Yes	No	Yes	tongue	16 HR-HPV
49	M	neoplasia	verrucous carcinoma	Yes	Yes	No	lip	16 HR-HPV
31	F	neoplasia	oral squamous cell carcinoma	No	Yes	No	buccal mucous	16 HR-HPV
71	F	neoplasia	oral squamous cell carcinoma	Yes	Yes	Yes	Retromolar Trigone	16 HR-HPV
89	M	neoplasia	verrucous carcinoma	Yes	No	No	buccal mucous	26 HR-HPV
35	F	neoplasia	carcinoma in situ	Yes	Yes	No	tongue	52 HR-HPV
60	M	neoplasia	oral squamous cell carcinoma	No	Yes	Yes	tongue	52 HR-HPV

Table 2. Statistics correlation between HPV and the different variables

	HPV			p <0.05 P value (<0.05)	OR 95 % C I	
	POS (%)	NEG (%)	β			
TOBACCO SMOKER				4.36	0.037	2.82 0.94-8.58
YES	12 (52.2)	11 (47.8)				
NO	17 (27.9)	44 (72.1)				
ALCOHOL DRINKER				4.48	0.034	3.09 0.94-10.34
YES	10 (55.6)	8 (44.4)				
NO	19 (28.8)	47 (71.2)				
GENDER				8.28	0.004	3.91 1.37-11.35
MALE	19 (51.4)	18 (48.6)				
FEMALE	10 (21.3)	37 (78.7)				
AGE				0.05	0.830	0.91 0.33-2.46
>50	13 (61.5)	26 (38.5)				
≤50	16 (46.4)	29 (53.6)				

Table 3. Statistics correlation between: lesions, cytology and localization of lesion vs HPV and conventional risk factors

	HPV			ALCOHOL DRINKER			TOBACCO SMOKER		
	POS (%)	NEG (%)	p (<0.05)	POS (%)	NEG (%)	p (<0.05)	POS (%)	NEG (%)	p (<0.05)
LESION			0.000			0.428			0.535
YES	29 (53.7)	25 (46.3)		13 (24.1)	41 (75.9)		16 (29.6)	38 (70.4)	
NO	0 (100.0)	30(100.0)		5 (16.7)	25 (83.3)		7 (23.3)	23 (76.7)	
BENIG LESION			0.020			0.887			0.594
YES	8 (88.9)	1 (11.1)		2 (22.2)	7 (77.8)		2 (22.2)	7 (77.8)	
NO	21 (46.7)	24 (53.3)		11 (24.4)	34 (75.6)		14 (31.1)	31 (68.9)	
PML			0.050			0.057			0.006
YES	12 (41.4)	17 (58.6)		4 (13.8)	25 (86.2)		4 (13.8)	25 (86.2)	
NO	17 (68.0)	8 (32.0)		9 (36.0)	16 (64.0)		12 (48.0)	13 (52.0)	
NEOPLASIA			0.808			0.028			0.001
YES	9 (56.2)	7 (43.8)		7 (43.8)	9 (56.2)		10 (62.5)	6 (37.5)	
NO	20 (52.6)	18 (47.4)		6 (15.8)	32 (84.2)		6 (15.8)	32 (84.2)	
CYTOLOGY									
GRADE II			0.023			0.152			0.191
YES	13 (76.5)	4 (23.5)		2 (11.8)	15 (88.2)		3 (17.6)	14 (82.4)	
NO	16 (43.2)	21 (56.8)		11 (29.7)	26 (70.3)		13 (35.1)	24 (64.9)	
GRADE III			0.027			0.637			0.171
YES	11 (39.3)	17 (60.7)		6 (21.4)	22 (78.6)		6 (21.4)	22 (78.6)	
NO	18 (69.2)	8 (30.8)		7 (26.9)	19 (73.1)		10 (38.5)	16 (61.5)	
GRADE IV and V			0.903			0.015			0.001
YES	5 (55.6)	4 (44.4)		5 (55.6)	4 (44.4)		7 (77.8)	2 (22.2)	
NO	24 (53.3)	21 (46.7)		8 (17.8)	37 (82.2)		9 (20.5)	36 (79.5)	
LOCALIZATION OF LESION									
TONGUE			0.050			0.531			0.121
YES	12 (41.4)	17 (58.6)		6 (20.7)	23 (79.3)		6 (20.7)	23 (79.3)	
NO	17 (68.0)	8 (32.0)		7 (28.0)	18 (72.0)		10 (40.0)	15 (60.0)	
BUCCAL MUCOUS			0.030			0.788			0.562
YES	11 (78.6)	3 (21.4)		3 (21.4)	11 (78.6)		5 (35.7)	9 (64.3)	
NO	18 (45.0)	22 (55.0)		10 (25.0)	30 (75.0)		11 (27.5)	29 (72.5)	

Due to the low number of cytological samples grades IV and V, both groups were gathered for the analysis

Table 3 shows a significant association between the presence of HPV and patients with lesions ($\beta= 24.61$, $p = 0.000$), benign lesions ($\beta= 5.38$, $p = 0.020$, $OR= 9.14$, $0.99-211.36$), cytological grade II ($\beta= 5.17$, $p = 0.023$, $OR= 4.27$, $1.01-19.31$) and location of the lesion on buccal mucosae ($\beta=4.70$, $p = 0.030$, $OR= 4.48$, $0.94-24.14$).

Likewise, we observed a significant association between alcohol drinkers with neoplasias ($\beta= 4.82$, $p = 0.028$, $OR= 4.15$, $0.93-19.17$) and with cytological grades IV and V ($\beta= 5.86$, $p = 0.015$, $OR= 5.78$, $1.02-34.75$). We observed a negative association between the presence of HPV and cytological grade III ($\beta= 4.86$, $p = 0.027$ $OR= 0.29$,

0.08-1.02). Tobacco smoking habit showed a significant association with neoplasias ($\beta= 11.78$, $p = 0.001$ OR= 8.89, 1.97-43.27) and with cytological grades IV and V ($\beta= 12.01$, $p = 0.001$, OR= 14.00, 2.07-119.84). We observed a negative association between tobacco smoking habit and the occurrence of lesions classified as PML ($\beta= 7.53$, $p = 0.006$, OR= 0.17, 0.04-0.75).

Males showed a significant association with the finding of lesions ($\beta= 5.72$, $p = 0.017$, OR= 3.19, 1.10-9.48) and with benign lesions ($\beta= 5.38$, $p = 0.020$, OR=9.14, 0.99-211.36). Age showed no significant association with any clinical histopathological variable (Data not included in Table 3).

4. Discussion

Understanding the role of HPV in oral carcinogenesis is hard due to the different frequencies of HPV infection in potentially malignant lesions and in oral cancer[23,24].

The percentage of HPV detection in our study was similar to data reported in other publications[25,26], likewise, no significant differences were observed between the detection of the different genotypes, HR-HPV and LR-HPV (14 vs 15 respectively). However, we can observe that the high-risk types were found mainly in malign lesions, while the low risk types were detected mainly in benign lesions. This becomes evident when analyzing the data shown in Figure 1, where is observed that HR-HPV was detected in 77.8% of cases of neoplasia (HPV+), while that LR-HPV was detected in only 22.2% of patients in this same group. An inverse situation is observed in the group of patients with benign lesions, showing clearly that HR-HPV is associated with increased severity of the injury.

Like in previous publications, HPV-16 was the HR-HPV genotype most frequently found[27,28] and 52 HR-HPV the second, but unlike other studies, genotype 18 was not detected at all in our work[28].

The presence of oral trauma and/or candida was included within the clinical histopathological variables analyzed; we did not find a significant association with HPV, history of alcohol intake, tobacco smoking, gender or age. Also, like other authors, our study did not show significant associations between HPV and patient's age[5,29].

Unlike other authors who have reported higher rates of HPV detection in patients without history of tobacco or alcohol intake[30,31], our study showed that patients who smoked tobacco or drank alcohol were 3 times more likely to have infections by HPV (Table 2).

Significant associations of HPV with cytological grade II, and tobacco and alcohol with cytological grades IV and V was in agreement with the finding of these risk factors for benign and malignant lesions, respectively. (Table 3).

On other hand, we observed a negative association between HPV with cytological grade III and tobacco smokers with PML, both with OR less than 1 (0.29 and 0.17) indicated that cytological grade III and PML are less frequent

in those patients with HPV infection and tobacco smoking history respectively[32].

As in other publications, the more frequent location of the lesions was the tongue[25,33] accomplishing 53.70% of the cases, followed by 25.93% in the buccal mucosae. In the statistical analysis, both tongue and buccal mucous showed no significant association with other groups (benign lesions, PML and neoplasias). The lesions of the buccal mucous presented significant association with HPV, which is in agreement with data previously reported by other authors on the basis of the epitheliotropic nature of the virus (table3) [11,12].

Our results showed that tobacco smoking and alcohol drinking have more influence than HPV on the development of neoplastic lesions. However the 56.2% of the neoplasias were positive for HPV (Table 3), and the percentage of HR-HPV detection increases with the severity of the lesions, suggesting the possible involvement of HPV in the malignant processes (Figure 1). Perhaps, may be that HR-HPV alone is not able to significantly increase the risk of developing malignant lesions of the oral cavity, but when in conjunction with other factors such as tobacco and alcohol, it develops an important role in developing these lesions.

Many studies have confirmed that HPV is the causative agent of cervical cancer, but its role as etiologic agent of oral cancer needs to be further studied to achieve accurate conclusions.

Funding

This study was supported by grants from the Fundación Roemmers subsidio 2012-2014 and Ministerio de Ciencia y Técnica de la Provincia de Córdoba-Argentina. PIO-MincytCba. 2011-2013. Ref. Res. [MINCYT Cba. N° 170/2011].

REFERENCES

- [1] Hirota SK, Migliari DA, Sugaya NN; 2006, Oral squamous cell carcinoma in a young patient - Case report and literature review. *An Bras Dermatol*; 81:251-4.
- [2] Llewellyn CD, Johnson NW, Warnakulasuriya KA (2004): Risk factors for oral cancer in newly diagnosed patients aged 45 years and younger: a case-control study in Southern England. *J Oral Pathol Med* 33:525-32.
- [3] Sasaki T, Moles DR, Imai Y, Speight PM (2005): Clinico-pathological features of squamous cell carcinoma of the oral cavity in patients <40 years of age. *J Oral Pathol Med* 34:129-33.
- [4] Gillison ML, Koch WM, Capone RB, Spafford M, Westra WH, Wu L, et al (2000). Evidence for a causal association between human papillomavirus and a subset of head and neck cancers. *J Natl. Cancer Inst* 92:709-20.

- [5] zur Hausen H. Papillomavirus and cancer (2002): from basic studies to clinical application. *Nat Rev Cancer* 2:342–50. S. M. Mitev and V. P. Veiko, *Laser Assisted Microtechnology*, 2nd ed., R. M. Osgood, Jr., Ed. Berlin, Germany: Springer-Verlag, 1998
- [6] Castellsague X, Quintana MJ, Martínez MC, Nieto A, Sánchez MJ, Juan A, et al. (2004). The role of type of tobacco and type of alcoholic beverage in oral carcinogenesis. *Int J Cancer* 108:741-9
- [7] Farshadpour F, Hordijk GJ, Koole R, Slootweg PJ (2007) Non-smoking and non-drinking patients with head and neck squamous cell carcinoma: a distinct population. *Oral Dis* 13:239-43.
- [8] Syrjanen K, Syrjanen S, Lamberg M, Pyrhönen S, Nuutinen J (1983) Morphological and immunohistochemical evidence suggesting human papillomavirus (HPV) involvement in oral squamous cell carcinogenesis. *Int J Oral Surg* 12:418–24
- [9] Erdmann J (2003). Recent studies attempt to clarify relationship between oral cancer and human papillomavirus. *J Natl Cancer Inst* 95:638–9
- [10] Nair S, Pillai M (2005) Human papillomavirus and disease mechanisms: relevance to oral and cervical cancers. *Oral Dis* 11:350–9.
- [11] Bernard HU (2005) The clinical importance of the nomenclature, evolution and taxonomy of the human papillomaviruses. *J Clin Virol* 32S:S1–S5
- [12] de Villiers EM, Fauquet C, Broker TR, Bernard HU, zur Hausen H (2004). Classification of papillomaviruses. *Virology* 20;324(1):17-27.
- [13] Bosh FX, Lorincz A, Muñoz N, Meijer CJ, Shah KV (2002). The causal relation between human papillomavirus and cervical cancer. *J Clin Pathol* 55:244–65.
- [14] Muñoz N, Bosch FX, de Sanjose S, Herrero R, Castellsagué X, Shah KV, et al (2003) Epidemiologic classification of human papillomavirus types associated with cervical cancer. *N Engl J Med* 348:518–27
- [15] Thompson IO, van der Bijl P, van Wyk CW, van Eyk AD (2001) A comparative light-microscopic, electron-microscopic and chemical study of human vaginal and buccal epithelium. *Arch Oral Biol* 46:1091–8.
- [16] Boy S, Van Rensburg EJ, Engelbrecht S, Dreyer L, van Heerden M, van Heerden W (2006). HPV detection in primary intraoral squamous cell carcinomas - commensal, a etiological agent or contamination? *J Oral Pathol Med*. 2006; 35:86–90.
- [17] Gillison ML, Broutian T, Pickard RK, Tong ZY, Xiao W, Kahle L, et al (2012). Prevalence of Oral HPV Infection in the United States, 2009-2010. *JAMA* 307(7):693-703
- [18] Warnakulasuriya S, Johnson NW, van der Waal I (2007). Nomenclature and classification of potentially malignant disorders of the oral mucosa. *J Oral Pathol Med* 36(10): 575-80
- [19] Herrero R, Castellsague X, Pawlita M, Lissowska J, Kee F, Balaram P, et al (2003). IARC Multicenter Oral Cancer Study Group: Human papillomavirus and oral cancer. The International Agency for Research on Cancer Multicenter Study. *J Natl Cancer Inst* 95:1772-83
- [20] Bernard HU, Chan SY, Manos MM, Ong CK, Villa LL, Delius H, et al (1994). Identification and assessment of known and novel human papillomaviruses by polymerase chain reaction amplification, restriction fragment length polymorphisms, nucleotide sequence, and phylogenetic algorithms. *J Infect Dis* 170:1077-85
- [21] De Roda Husman AM, Walboomers JM, Meijer CJ, Snijders PJ (1995). The use of general primers GP5 and GP6 elongated at their 3' ends with adjacent highly conserved sequences improves human papillomavirus detection by PCR. *J Gen Virol* 76:1057-62
- [22] InfoStat[computer program] versión 2011. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina. Available from: <http://www.infostat.com.ar>
- [23] Miller CS, White DK (1996) Human papillomavirus expression in oral mucosa, premalignant conditions and squamous cell carcinoma: a retrospective review of the literature. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 82:57–68.
- [24] Uobe K, Masuno K, Fang Y, Li LJ, Wen YM, Ueda Y, et al (2001) Detection of HPV in Japanese and Chinese oral carcinomas by in situ PCR. *Oral Oncol* 37:146–52.
- [25] Anaya-Saavedra G, Ramírez-Amador V, Irigoyen-Camacho ME, García-Cuellar CM, Guido-Jiménez M, Méndez-Martínez R, et al (2008). High association of human papillomavirus infection with oral cancer: a case-control study. *Arch Med Res* 39(2):189-97.
- [26] Tinoco JA, Silva AF, Oliveira CA, Rapoport A, Fava AS, Souza RP (2004). Human papillomavirus (HPV) infection and its relation with squamous cell carcinoma of the mouth and oropharynx. *Rev Assoc Med Bras* 50(3):252-6
- [27] Kreimer AR, Clifford GM, Boyle P, Franceschi S (2005) Human papillomavirus types in head and neck squamous cell carcinomas worldwide: a systematic review. *Cancer Epidemiol Biomarkers Prev* 14:467–75.
- [28] Termine N, Panzarella V, Falaschini S, Russo A, Matranga D, Lo Muzio L, et al (2008). HPV in oral squamous cell carcinoma vs head and neck squamous cell carcinoma biopsies: a meta-analysis (1988-2007). *Ann Oncol* 19(10): 1681-90
- [29] Badaracco G, Venuti A, Morillo R, Muller A, Marcante ML (2000). Human papillomavirus in head and neck carcinomas: Prevalence, physical status and relationship with clinical / pathological parameters. *Anticancer Res* 20:1301-6
- [30] Hafkamp HC, Manni JJ and Speel EJ (2002). Role of human papillomavirus in the development of head and neck squamous cell carcinomas. *Acta Otorinol* 124:520-6.
- [31] Ringstrom E, Peters E, Hasegawa M, Posner M, Liu M, Kelsey KT (2002). Human papillomavirus type 16 and squamous cell carcinoma of the head and neck. *Clin Cancer Res* 6:3187-92.
- [32] Sócrates Aedo M, Pavlov SD, Clavero FC (2010). Riesgo relativo y Odds ratio ¿Qué son y cómo se interpretan? *Rev. Obstet. Ginecol* 5 (1):51-4
- [33] Falaki F, Dalirsani Z, Pakfetrat A, Falaki A, Saghravanian N, Nosratzahi T, et al (2011) Clinical and histopathological analysis of oral squamous cell carcinoma of young patients in Mashhad, Iran: a retrospective study and review of literature. *Med Oral Patol Oral Cir Bucal* 16(4):e473-71.