

Mechanisms of Groundwater Pollutants Transport in Tulkarm Area / Palestine

Saed Khayat¹, Amer Marei^{2*}, Basel Natsheh¹, Nawaf Abu-Khalaf¹

¹Palestine Technical University (PTUK), Tulkarm, Palestine

²Environmental Research Lab., Al-Quds University Abu Dis, East Jerusalem, Palestine

Abstract This study aims at investigating the mechanisms of pollutants transport based on geological formations. The study classifies two sub-aquifers groups. The first group are wells which dogged within Senonian Abu Dis formation, and are mostly used extensively for municipal purposes. The second group are wells dogged with the most top Jerusalem-Hebron formation of Turonian age and mostly used frequently for irrigation purposes. The first group shows a high deteriorated water quality with ionic molar ratios bears the ratios from Wastewater stagnant in adjacent Zomar stream. The stream bed lays directly over Jerusalem formation which promotes the pollutant transport directly to the uptake zone. However, the effect of such stagnant become less in dilution time of winter. Septic tanks surrounded these wells are constructed within the top 5 meters of Senonian formation, which mean that the leakage from septic tanks is not that ease to significantly affect the groundwater quality. The second group are mostly subjected to direct rainfall infiltration this can be indicated through the higher Sodium values which are related to the ion exchange through limestone profile. The results emphasize the need for further procedures to mitigate the pollutants plume migration from Wadi to the surrounded environment.

Keywords Groundwater, Pollution, Palestine, Tulkarm, Karst, Stagnant Water, Transoundary Pollution

1. Introduction

Water is one of the most valuable natural resources in the Middle East. The combination of multiple factors, e.g. population growth, economic and agricultural development, and an arid climate, result in over exploitation of the water resources in the region. The continuous high water demand leads to rapid degradation of the quality of fresh water resources as a result of the salinisation and contamination processes[1-3]. The lack of sufficient water, combined with rapid water quality deterioration, presents a serious challenge to the people in the region. In order to manage and share the water resources under conditions of accelerating degradation, it is crucial to understand the origin and mechanisms of the contamination process.

The salinity that threatens the fresh water resources is derived from different sources, both natural and anthropogenic. In general, over exploitation of fresh aquifers results in a rapid decrease of water level, which then triggers lateral as well as upwelling of deep saline water from adjacent waters from aquifers. Consequently, the overexploited aquifers become saline due to mixing with salinewaters[1-4].

Tulkarm area is located in the north western part of the West Bank. The water resources in the district is limited and restricted only to the shallow aquifer that is subjected to various sources of pollutant ion.

Groundwater in Tulkarm City is being utilized through 10 domestic and 53 irrigation wells with total discharge of 21 MCM per year[5]. Most of the wastewater from the local communities to the east of Tulkarm drained through long stream that extended from Nablus city to the east across Tulkarm city to the Midetranean Sea. The wastewater stream part that pass from Nablus through Tulkarm is called Wadi Zomar. The main objective of this study is to investigate the role of different geological formations in facilitating the transport of the pollutants from Wadi Zomar and different anthropogenic sources surrounding Tulkarm city in Palestine.

2. Study Area

Tulkarm area is located in the north western part of the West Bank - Palestine (Figure 1). The total area of Tulkarm is about 246 km² and its current population is estimated at 166,832 people, representing 12.4% of the total population of the West Bank[6]. Tulkarm lies on the western slopes of the West Bank, which are characterized by gentle slopes. The elevation in the city ranges between 50 to 180 meters above sea level.

* Corresponding author:

marei.amer@gmail.com (Amer Marei)

Published online at <http://journal.sapub.org/re>

Copyright © 2012 Scientific & Academic Publishing. All Rights Reserved




Figure 1. Study Area of Tulkarm, sampled wells are shown with the different geological formations

The climate in Tulkarm is similar to that of the Mediterranean type, which is characterized by long, hot, dry summers and short, cool, rainy winters. Tulkarm is distinguished by the moderating effect the sea breeze has on its climate. Rainfall is limited to the winter and spring months, mostly between November and March while summers are completely dry. The average annual precipitation rate in Tulkarm is about 640 mm[7].

Tulkarm is supplied with water from municipal well no. 1 (Fig.1) with production capacity of 180 m³ / hour, well no. 2 with production capacity of 100 m³/hour, well no. 3 with production capacity of 65 m³/hour and Shwekeh well with production capacity of 110 m³/hour. In summer time, where water consumption increases and production capacity decreases, the Municipality of Tulkarm makes use of few agricultural wells like Al Taffal Well, Hannun Well, Abdel Razeq Well and Al Safarini Well to sublement the increase in water demand. The average water table in the Municipality wells is 90 meters deep. Water consumption is 85 L/Capita/Day.

There are many wadis (streams) located in the district. The main sewage wadi in the region is called Wadi Zomar. The hydrographic catchments area of the Zomar Valley is about 22 km long and 5 km wide[7]. Wadi Zomar exposed to major pollution by human activities, that causes serious health problems and other economic costs related to water treatment, remediation and locating a new water supply,

become evident[8]. The Wadi Zomar is used as drainage for the sewage from the towns of Nablus, Tulkarm and the villages located at the banks of the Wadi.

There are still more than 500 septic tanks/cesspools, which are rarely emptied. Water analysis in the Municipal well no. 1 made three years ago demonstrates pollution of ground water and presence of Fecal Colifrom.


Previous studies for the water quality in the study area shows that there is a big pollution problem with sewage especially for municipal wells which used for domestic purposes[9-16]. These studies try to connect the problem of groundwater pollution to the seepage from the septic tanks. Less were talking about the role of Wadi Zomar in surrounding environment pollution[17]. Moreover, the role of different geological formations in pollution transport still arbitrary.

2.1. Geology

The geological formations of the study area of Tulkarm range in age from Senonian to Quaternary. The stratigraphic section of the West Bank is present in (Fig.2)[18]. The district is mainly covered by sedimentary carbonate rocks such as limestone, dolomite, marl and chalk. The general geology of the Tulkarm area is represented in Figure 1. A brief description of the lithological formations encountered in the Tulkarm district is presented below[18-20].

Period	Age	Graphic Log	Typical Lithology	Formation (West Bank Terminology)	Sub-Formation	Group	Symbol	Formation (Israeli Terminology)	Hydro-stratigraphy	Typical Thickness (m)							
Quaternary	Holocene		Nari (surface crust) and alluvium Gravels and fan deposits	Alluvium			Qh-a	Alluvium	Local Aquifer	0 - 100							
	Pleistocene		Thinly laminated marl with gypsum bands and poorly sorted gravel and pebbles	Lisan			Qp-l	Lisan(Kurkar Group)	"Aquitard"	10 - 200							
Tertiary	Miocene Pliocene		Conglomerates, marl, chalk clay and limestone	Beida			Tmp-b	Saqiyeh Group	Local Aquifer	20 - 200							
	Eocene (Lower - Middle)		Nummulitic reefal Limestone Nummulitic bedded Limestone Nummulitic Limestone,Chalk Chalk ,Nummulitic Limestone	Jenin	Jenin 4 Jenin 3 Jenin 2 Jenin 1	Jenin	Te-j Te-j4 Te-j3 Te-j2 Te-j1	'Avedat Group	Aquifer	90 - 670							
Cretaceous	Paleocene		Marl,Chalk	Khan Al-Ahmar		Nablus	Ks-n	Mt.Scopus Group	Aquitard (Local Aquifer)	40 - 150							
	Maasirich-tian Danian		Chalk ,Marl	Wadi Al-Qilt						10 - 120							
	Campanian		Main Chert ,Phosphate	Abu Dis					0 - 450								
	Upper	Turonian		White Limestone ,stilloithes Limestone and Dolomite Yellow thin bedded Limestone	Jerusalem	Upper Middle Lower	Ramallah	Kc-j	Bina	Upper Aquifer	40 - 190						
		Cenomanian		Dolomite,soft	Bethlehem	Upper					Kc-b	Weradim Kefar Sha'ul	50 - 210				
	Lower	Albian		Karstic Dolomite	Hebron	Lower	(West Bank)	Kc-h	Amminadav	Lower Aquifer	65 - 160						
				Yellow marl Lime & Dolostone,Chalk,(Clay)	Yatta	Upper Lower					Kc-y	Moza Beit Meir	"Aquitard"	50 - 125			
		Aptian		Reefal Limestone	Upper	UBK2						Kobor	Ka-ubk	Kesalon Soreq	Lower Aquifer	10 - 20	
				Dolomite Limestone, interbedded with Marl	Beit Kahil	UBK1					Ka-lbk					Giv'at Ye'arim Kefira	80 - 130
				Dolomite	Lower	UBK2											Ka-lbk
			Karstic Limestone	Beit Kahil	UBK1	Ka-lbk					100 - 160						
Neocomian		Marl ,marly nodular Limestone	Qatana		Kurnub	Ka-q	Qatana	Aquitard	42								
		Marly Limestone and Limestone	Ein Qinya						Ka-eq	Ein Qinya	Local Aquifer	55					
		Shale	Tammun									Ka-t	Tammun	Aquitard	300+		
	Shale and Limestone	Ein Al-Assad		Ka-ea				20+									
Jurassic	Oxfordian		Marly Limestone,sandy					Nabi Sa'id		Kurnub	Ka-rs	Hatira	Aquifer	20+			
			Sandstone	Ramali		Kn-r								70+			
	Volcanics	Tayasir			Kn-t								35				
	Marl interbedded with chalky limestone	Maleh	Upper Maleh			Jo-um	'Arad Group		Aquitard				100 - 200				
	Dolomitic limestone, jointed and karstic	Maleh	Lower Maleh		Jo-lm							Aquifer	50 - 100				

Stratigraphic Section of the West Bank



LEGEND

	Dolomite		Mega/auna		Sandstone
	Limestone		Flint concretions		Volcanics
	Marl		Chalk		Relatively Permeable
	Conglomerate		Nari		Relatively Impermeable

Figure 2. Stratigraphical section of the West Bank (Source:[18])

Table 1. Hydrochemistry of the Pleistocene Wells and springs from Jericho-Palestine area in (mg/l)

Sample ID	X	Y	Well Depth (m)	Abstraction rate. (MCM/yr)	NO ₃	pHfield	Conductivity (µs/cm)	PO ₄	SO ₄	Cl	HCO ₃	Mg	Ca	Na	K
Munic. Well	152400	190920	110	0.432689276	64	7.01	1026	7.2	40	95.72	512.6	40.25	96.42	66.07	24.62
Fadilyeh Munic.	152480	191900	118	0.348840931	49	6.95	1104	1.7	39	106.35	512.6	40.44	98.22	70.85	23.15
Shufa Committee	155500	188500	151	0.066066556	28.3	7.1	741	0.13	20	42.54	524.8	38.54	90.15	33.06	2.16
Isma'el Itair	153750	188800	100	0.18636172	59.4	7.09	815	0.6	43	49.63	414.9	37.04	94.73	40.95	3.57
Izbeh Well 5 Munic	154451	191291	186	0.18636172	54.3	7.06	760	0.7	25	53.18	366.1	32.94	82.25	44.15	4.8
Izbeh Well 6 Munic	153459	190004	200	0.18636172	41	7.06	794	0.2	26	53.18	427.1	22.01	55.55	35.6	1.1
Muhammad Omar	152350	189000	74	0.119719417	23	7.12	667	0.13	16	42.54	451.5	33.15	73.46	35.21	2
Muhd ABED AL HALEEM	152500	187950	165	0.171508167	23.5	7.01	689	0.8	13	46.09	427.1	35.85	72.99	34.59	1.12
As'ad Taffal	154440	190880	136	0.153256042	37.4	7.07	650	0.26	16	42.54	378.3	30	65.99	38.61	1.21
Sadeeq Jamus	154770	190910	157	0.157739458	57.4	7.07	725	0.5	27	53.18	353.9	33.4	75.94	40.78	1.24
Hasan Khaleel	156800	190100	175	0.093099417	32.4	7.27	571	0.08	22	31.91	341.7	29.45	60.62	28.99	2.49
Zubaydah Al Sa'eed	155730	191000	154	0.141912708	45	7.39	740	0.16	21	67.36	366.1	34.86	71.2	45.85	4.91
Rafeeq Hamdallah	154540	192040	90	0.132879792	53	7.03	889	0.06	23	81.54	427.1	35.15	88.8	63.19	4.36
Hafedh Alhamdallah	153470	191600	80	0.113321625	68	6.84	1062	0.4	21	92.17	549.2	40.59	110.87	62.77	9.16
Municipality Shweikeh	152480	191900	118	0.348840931	45.9	6.93	1033	0.1	36	102.81	500.4	41.96	90.47	69.78	15.49
Almisnawi Munic,	153259	193780	125	0.348840931	57	6.92	947	2.5	23	85.08	475.9	39.56	88.1	60.82	16.7
Ali Abu Saleh	152400	193040	132	0.169933833	32.7	6.95	993	0.06	31	106.34	488.2	40.73	91.4	71.17	8.03
Wadi Zomar Wastewater	153323	193241	-----	52 *	189	6.23	6234	31.2	141	273	1024	87	214	185	53

*For the whole tributaries according the World Bank report on Water status in the West Bank, 2011

2.1.1. Upper Cenomanian

The Upper Cenomanian formation (also known as the Bethlehem formation) consists of limestone, dolomite with some chalk, and marl. Outcrops are found mainly on the flank of the A'nabta anticline. The dolomite forms a rugged morphology on gentle slopes.

2.1.2. Turonian

The Turonian formation (also known as the Jerusalem formation) consists of a series of massive, thick- to- thin bedded limestone to dolomitic limestone and dolomites with a thickness of approximately 70-130 m. The lower part of the Turonian formation consists mainly of limestone and dolomite with marl and some chalk, making it sometimes difficult to be distinguished from the underlying Bethlehem formation. Towards the top of this formation, chalk beds with occasional chert bands are common, and the formation is transitional to the overlying chalk facies. The Turonian formation has a well-developed karst feature and is commonly used as a building stone. It is exposed in the A'nabta anticline and is considered a very good aquifer.

2.1.3. Senonian

The Senonian formation (also known as Abu Dis formation) is mainly made up of Cretaceous Rocks, which are composed of chalk, ranging in age from the Coniacian at the bottom to Paleocene at the top. Outcrops exist in the A'nabat anticline and on the western limb of Nablus-Beit Qad syncline. In the Tulkarm city area, the chalk is thin and consists of marly base and passes upwards through bedded and crystalline limestone that has few marl partings.

2.1.4. Quaternary

Quaternary rocks are divided into the following formations:

a. Lisan Formation: these recent sediments are mainly composed of alluvium consisting of bedded limestone, chert and clay. The thickness of these sediments varies from one place to another. The rock fragments comprising the deposits are mainly derived from rock formations adjacent to the wadi area.

b. Nari Formation: it occurs mainly in high rainfall areas where carbonate rocks are dissolved by percolating water. It forms a thin coating over the limestone.

2.2. Hydrogeology

The Tulkarm district is underlined hydro-geologically by the northern part of the western groundwater basin. It is located in the semi-coastal region. Groundwater flow is directed towards the west and northwest.

2.2.1. Aquifer System

The major aquifers in the area are the Shallow Aquifer and the Upper Cenomanian- Turonian complexes. The Abu Dis acts as an aquitard hence, forming a water barrier. Since

the Upper Aquifer is the predominant aquifer in the region, it is described in full below [19,20].

2.2.1.1. The Upper Aquifer System

The Upper Aquifer is represented by the formations of the age Turonian (Jerusalem formation) and Cenomanian (Bethlehem and Hebron formations).

2.2.1.2. Turonian Aquifer

The Turonian aquifer is part of the Upper Aquifer but can be classified as a distinct local aquifer if the formation beneath it acts as an aquitard as is the case in some areas in the eastern and southern parts of the West Bank. The Turonian aquifer is considered a fairly good aquifer especially where the saturation thickness is in tens of meters. This aquifer is of good thickness and extent in the Tulkarm area (approximately 130 m thick). The water quality of this aquifer is generally good but in some areas there is evidence of deterioration because of sewage and agro-chemical pollution.

2.2.1.3. Cenomanian Aquifer

The formations of this aquifer are the Bethlehem and Hebron. The aquifer is an important regional source of water supply for domestic use. The Cenomanian Aquifer has high recharge values. It is heavily exploited in the areas near Tulkarm and Qalqilya because the aquifer is at a shallow depth in this area. The depth to water is rarely more than 200 m below ground surface.

3. Data Collection and Method

The original aim of groundwater sampling was based on providing a supplement analyses to investigate the gradient of pollution for the aquifer system, according to spatial distribution and the effect of the pollution plume surrounded the wastewater stream banks.

For this purpose, 17 samples from springs and wells, distributed in different geological outcrops, were collected.

The samples were collected in December 2011 (wet season), sampling localities are shown in Fig. 1. Other samples for Nitrate and potassium were obtained 4 months earlier by the end of the dry season.

Onsite measurements for physicochemical parameters (pH, temperature, m-value and redox potential) were done. The samples for anions were preserved using HgCl₂ for prolonged preservation of nitrate. For accurate chloride measurement, the chloride concentration was measured out from the water samples of the cations [3]. Due to the fact that samples were collected in the agricultural season, all the sampled wells had been pumping for a long period of time. Hence, no further purging was needed prior to sampling and the samples were collected immediately. Samples for (Cations and Anions) were collected in 250 ml polyethylene bottles.

Major anions (NO_3 , SO_4 , PO_4 and Cl) were analyzed using HP liquid chromatography. Concentrations of the major cations (Ca, Mg, Na, and K) were determined by ICP-OES. Alkalinity and HCO_3 analysis was measured onsite by titration[21,22].

All chemical analyses were carried out at the Environmental Research Lab. at Al-Quds University – Abu Dis.

4. Results

The chemical data are shown in Table 1. The major cations and anions concentrations of the groundwater from the sampled wells in the study area are plotted on a piper diagram (Figure 3).

The water type is normal earth alkaline water with prevailing bicarbonate for one group and prevailing chloride in other group. The water type varied from

Ca-Mg- HCO_3 which shows less susceptibility to pollution to those have Ca-Mg-Na- HCO_3 -Cl for those wells with high pollutants, the last type bears the sample from Wadi Zomar wastewater (Figure 3). However, both groundwater types reflect dolomite bearing rock from Turonian and sometimes the deeper cenomanian formations.

In general the results shows different NO_3 values in all samples that varied from 64 mg/L in winter season and 81 mg/L in summer season for municipal wells, to 23 mg/L in Muhammad Omar agricultural well. In general most of the nitrate vales exceed the WHO standard for safe drinking water which is 50mg/L. While other anthropogenic related ions show variable values which exceed in many cases the WHO standard for drinking water and follow the surrounded activities and the characteristics of the aquifer geology as it will be described in further details later on in the following section.




Figure 3. Durov Plot suggests the trend of water from fresh Ca-Mg- CO_3 water in some wells NA-Mg-Cl water in others that are bearing the Wadi Zomar value

5. Discussion

In general, many previous studies for the water quality in the study area show that there is a big pollution problem with sewage especially for municipal wells which used for domestic purposes. These studies mostly refer the problem of groundwater pollution in the area to the seepage from the septic tanks[9-16]. This study shows that these previous results are partially true. In this study, we tried to make further geological investigation to build a conceptual model for pollutants transport, whether it was from septic tanks or from Wadi Zomar. There are some other studies taking

Wadi Zomar as environmental hazards for the surrounding communities from Nablus to the trans-boundary area across green line to the Israeli areas[15,17]. However, the question about the mechanism of pollutants transport from the Zomar stream to the surrounded environment still open. The hydrochemical data from the groundwater in sampled wells surrounding Tulkarm city, show two dominant groups that have different magnitude in pollution and pollution type. In general the high pollution was recognized in the wells that have higher abstraction rate, mainly those wells that used at daily basis for domestic purposes.




Figure 4. Na vs. Chloride and Na/Cl molar ratios in the samples wells shows 2 major groups, one with high Na and Cl the same as in Wadi Zomar, and the other with relatively lower values




Figure 5. K/NO₃ molar ratio versus PO₄ in mmol/L, classified the samples into Wadi Zomar affected group, septic tanks affected group and wells with mixing effects

According to piper diagram those municipal wells group characterized by higher Na and Cl values that are more closed to those values taken from Wadi Zomar stream (Figure 3). The other group has less Cl content and more Na than chloride which might be due to ionic exchange during the process of recharge (Figure 4). In this context, it is worthy to mention that most of the wells exposed to normal softening are covered with Turonian outcrop which promote the direct infiltration through surface, unlike the other group that is mostly covered by impervious chalky Senonian formation.

The hydrochemical data classified the sampled wells into two groups which are subjected mainly to three pollutants sources (Figures 3, 4, 5). In general the pollutants sources varied according to the geological outcrop, and surrounded activities. These sources can be summarized as follow:

5.1. Wadi Zomar

Wadi Zomar is considered one of the largest wastewater streams that hold the wastewater from the surrounded communities and extend from Nablus city in the east and cross the green-line boundary to Israeli settlements across Tulkarm area. The problem of Wadi Zomar stream varied spatially and temporally, according to the season and the distance from the wastewater inlet and the slope[17]. Two geological profiles in the study area were created depending on the wells lithology (Figures 6, 7). The profiles show that most of the wells in the study area are drained from the Turonian- Cenomanian layers (Jerusalem-Hebron formation) (Figures 2). The Turonian Jerusalem formation is exposed in the area surrounded the Zomar stream, while the Cenomanian Hebron formation is found on the flank of the A'nabta anticline upper part of the stream (Figure 1, 2, 6, 7). The direct seepage from Hebron formation allowed the sewage to mix directly through with fresh recharge water and leaks to the zone of uptake (Figure 7).




Figure 6. Concentration of Potassium and Nitrate (in mg/L) in Summer and Winter season, show an increase in concentration in the dry season




Figure 7. A-B geological cross section from the North East to the South West of the study area

Two main factors increase the problem of wastewater leakage, these factors can be summarized as the increasing the abstraction pressure in the nearby municipal wells, and formation of wastewater stagnant, where the slope of the wastewater stream become more gentle and semi-straight in the study area, allowing the formation such wastewater stagnant which promote the wastewater to infiltrate slowly to the groundwater. This can be indicted through the variation of Potassium and nitrate values from winter to summer season (Figure 8), where in winter the stream receive additional runoff amount from the precipitation that prevent the formation of wastewater stagnant across the flow path.




Figure 8. C-D geological cross section from the East to the West of the study area

In general, both conditions favoured an environment where the pollution plume surrounded the wadi stream pulls in toward the zone of uptake through Jerusalem Hebron formation (Figures 6, 7). The most highly abstracted wells group (containing Municipality 1, Municipal Fadilya 2, Shweikeh Municipality well 3, and Misnawi Municipality well, in addition to two wells lay directly near the Zomar stream) are characterize by higher amount of Cl, PO₄, and K. The values for these elements are bearing the values from

samples obtained from wadi Zomar wastewater (Figure 5).

5.2. Septic Tanks

As mentioned above, several studies and reports connected the problem of high nitrate in the groundwater from the study area to the septic tanks distributed over the study area. Septic tanks, naturally, play an important role in the pollution process but the complete story is not only related to the septic tanks and it is not definitely the main cause of groundwater pollution as this study pinpoint. The data of this study suggest more important role for Wadi Zomar in the pollution process rather than the effect of cesspits. As mentioned above, this suggested role for the Wadi Zomar is clear in the wells of high abstraction rate and which are dogged in the impervious layers of Senonian (Figure 1). Normally septic tanks are present in the top 5 m below surface; therefore, the evidence of leakage from septic tanks through the Chalky Senonian layer is very low. However, a significant evidence of pollution from septic tanks was noticed in the wells dogged in the Turonian layer, these wells shows higher nitrate over Potassium values with relatively low PO_4 which found higher in wastewater from Wadi Zomar (Figure 5). These wells (dogged in Turonian) also show surplus Na in most cases which might indicate softening during the direct recharge through Jerusalem formation or might due to additional source of Na from Septic tanks with Na/Cl molar ratio higher than this ratio taken from Wadi stream (Figure 4).

5.3. Agricultural Activities and Irrigation Feedback

The protected agriculture is widely common all over Tulkarm area, where the drip irrigation practices are mostly used; however, some farmers are still using direct water application for irrigation. Moreover, there are a lot of agrochemicals and fertilizers applied in considerable amount in the study area[10]. In general, there are many indicators about the presence of irrigation feedback to the groundwater. Most of the farmers use Potassium nitrate and natural manure as fertilizers. This resulted in a surplus amount of potassium and nitrate in the groundwater. However, the data shows a wide range of both potassium and nitrates in the groundwater samples. In this context, it was difficult to differentiate between those values that are coming from the pollution plume of the Wadi Zomar, septic tanks or the agricultural feedback. Figure 5 shows that there are some wells (s. a. Rafeeq Hamadallah, Izbeh wells, and Ismael Itair) which are less affected by Zomar sewage plume, where the contribution of agricultural pollutants are clearer with higher potassium and relatively low PO_4 . This means that most of the wells are subjected to such kind of pollution, but the presences of more effective sources are dominated over the agricultural sources.

6. Conclusions

The problem of groundwater pollution in the upper

aquifer of Turonian Cenomanian age is the most common problem for the groundwater in the study area. Previous studies connected the source of pollution by the presence of common used septic tanks, agricultural activities and to limited extend the wastewater discharge in Wadi Zomar. Unlike other studies, this article classifies the pollution sources and the predominance effect of each source on each wells group. The study connects the different mechanisms of pollutants transport to the different geological pollution that characterize the surrounded aquifer media. In general, the study classifies two wells groups according to its location within geological outcrops. These are wells which dogged with the most top Senonian Abu Dis formation, which are mostly municipal with high daily abstraction rates, and the wells dogged with the most top Turonian Jerusalem formation and mostly used frequently for irrigation purposes. The first well group show a high deteriorated water quality that mostly related to the pollutant transport from Wastewater stagnant in Wadi stream. The stream bed lays directly over Jerusalem formation which promote the pollutant transport directly to the uptake zone of these wells (especially municipal one with high abstraction rate), and overcome any other pollution sources that might affect the quality of groundwater in these wells. Moreover, the septic tanks surrounded these wells are present in the most top 5 meters of impervious Senonian Abu Dis formation, which mean that the pollution from septic tanks has less effect on the groundwater from it. Moreover, the flatten slope of the Wadi stream surrounding Tulkarm enhance the formation of wastewater stagnant that infiltrate slowly through the alluvial deposits to Jerusalem formation and finally to the surrounded aquifer. The formation of such wastewater stagnant is mostly predominant in Sumer season with the absence of sufficient waste dilution or precipitation runoff that wash out the Wadi flow.

On the other hand, the second wells group shows higher sodium values which are related mostly to the ion exchange or softening that occurs with direct infiltration through Jerusalem formation cover the top of these wells locations. However, this group show less values of pollutant, and accordingly the effect of other pollution sources was easily noticeable. This group characterize by more effect from the septic tanks, then the pollutants from irrigation feedback.

7. Recommendations

The process of pollutants plume migration from Wadi Zomar through the geological formations to the adjacent uptake zone, can be controlled through lining the wastewater stream with suitable impervious material that prevent the leakage from the Wadi along the Stream path.

The daily heavy abstraction rate from the adjacent municipal wells must be regulated and the change in water quality should be monitored with respect to fluctuation of water table.

The effect of wastewater stagnant formation especially in summer season must be mitigated by injecting the stream with a suitable amount of treated wastewater that accelerate the stream flow and wash these stagnant out along wadi path.

ACKNOWLEDGEMENTS

We express our deep gratitude for Tulkarm Municipality for their cooperation in completing this work. In this regard we express our deep thank here to Eng. Rayek Hamad and Mr. Nidal Alshen from Water department in Municipality for all assistant and cooperation in field and providing data. The authors also would like to thank the Palestine Technical University-Kadoorie for providing financial support to publish this work.

REFERENCES

- [1] Vengosh, A., Rosenthal, E., 1994. Saline groundwater in Israel: its bearing on the water crisis in the country. *Journal of Hydrology* 156, 389-430.
- [2] Salameh E., 2002. Sources of water salinities in the Jordan Valley area/Jordan. *Acta Hydrochim Hydrobiol* 29:329–362.
- [3] Khayat, S., Hötzl, H., Geyer, S., Ali, W. 2006. Hydrochemical investigation of water from the Pleistocene wells and springs, Jericho area, Palestine *Hydrogeology Journal* 14 (1-2), 192-202.
- [4] Marie, A, Vengosh, A., 2001. Sources of salinity in groundwater from Jericho area, Jordan valley. *J Ground Water* 39:240–248.
- [5] MOPIC (Ministry of Planning and International Cooperation), 1998. Sensitive Water Resources Recharge Areas in the West Bank Governorates. 2nd Ed., West Bank Palestine, Ramallah.
- [6] PCBS (Palestinian Central Bureau of Statistics), 2010. Demographic Survey of the West Bank and Gaza Strip. District Report Series. No. 1-9. PCBS; yearly updates.
- [7] OPTIMA, 2007. D11.2 the Zomar stream-Wadi Zomar Basin. Optimization for Sustainable Water Resources Management.
- [8] Gasana, J., Morin, J., Ndikuyeze, A., Kamoso, P., 2002. Impact of water supply and sanitation on diarrheal morbidity among young children in the socioeconomic and cultural context of Rwanda (Africa). *Environ. Res.*, 90: 76-88.
- [9] Rabbo, A., Scarpa, D., Qanam, J., Jaber Z. Q. A., Younger, P., 1999. Wells in the West Bank, Water Quality and Chemistry, Palestinian Hydrology Group, University of Newcastle upon Tyne and Bethlehem University, Jerusalem, West Bank.
- [10] Qare, A., 2009. Potential Leaching of Pesticides to Ground-Water: A Look At The Most Common Use Pesticides In Palestine,” *Proceedings of the 2nd International Conference on Water: Values and Rights, Ramallah, 13-15 April.*
- [11] Masri, M. N., Abu Jaish, A., 2007. Emergency water supply and conservation in rural area impoverished by the wall: A qualitative research study on the water situation in the Tulkarem and Jenin governorates. A report submitted to CARE International. Water and Environmental Studies Institute, An-Najah National University, Nablus, Palestine.
- [12] Zayed, O., Abadi, M., Rajoub, B., Moe, H., 2005. Groundwater Vulnerability of Palestine Drinking Water Sources in the West Bank. In: *Water Values and Rights*. Palestine Academy for Science and Technology. Palestine Academy Press. Palestine.
- [13] Shahab, S. J., 1997. Hydrochemical and environmental aspects of the upper aquifer system in Tulkarm-Qalqeelya area, Palestine. MSc thesis. University of Jordan. Jordan.
- [14] Scarpa, D., Rabbo, A., Quannam, Z., Abdul-Jaber, Q., 1998. Groundwater Pollution in Unconfined Aquifers in the Northern West Bank, Palestine. *Proceedings of the British Hydrological society International Conference*. Exeter, Hydrology in a Changing Environment. Vol. II. John Willey and Sons. Chichester.
- [15] Hind M., 2004. Investigation on Groundwater Pollution, Field investigation in special locations in North and West of the West Bank. *Palestinian Wastewater Engineers Group (PWEG)*. Unpublished proceeding report. September 2004.
- [16] Naser, O., 2011. Kinetics of Groundwater Nitrate (NO₃-1) In Soil Leachate In Tulkarem City Using Soil Columns. MSc. Thesis. An-Najah University. January 2011.
- [17] S. Saleh, Mimi, Z., Khayat S., 2011. Using Biological Indicators to Characterize the Natural Flow Regime in Wadi Zomar Stream/Palestine. *Asian Journal of Applied Sciences*, 4: 685-701.
- [18] Aliewi, A., Rajab, K., Imseih, N., Abu Sada, J., Freimuth, L., Saltman, A., 2008. Protecting Trans-boundary Groundwater Sources from Pollution: ‘Research, Training and Guidelines for Palestinian and Israeli Municipalities’. House of Water and Environment, Palestine, and Friends of the Earth- Middle East, Tel Aviv.
- [19] PWA (Palestinian Water Authority), 2001a. Boundaries of the Western Aquifer Basin and the Eocene Aquifer in the Northeastern Aquifer Basin. Working Report SUSMAQ-MOD # 6.1 V1.0. Palestine.
- [20] PWA (Palestinian Water Authority), 2001b. Data Review on the West Bank Aquifers. Working Report SUSMAQ-MOD # 02 V2.0. Sustainable Management of the West Bank and Gaza Aquifers. Water Resources and Planning Department. Palestinian Water Authority. Palestine.
- [21] Connolly, D. and Paull, B. 2001. Rapid determination of nitrate and nitrite in drinking water samples using ion-interaction liquid chromatography. *Analytica Chimica Acta*, 441: 53-62.
- [22] Jackson, PE. 2001. Determination of inorganic ions in drinking water by ion chromatography. *TrAC Trends in Analytical Chemistry*. 20: 320-329.