

Survival Analysis of Under-five Mortality and Its Associated Determinants in Nigeria: Evidence from a Survey Data

Anthony I. Wegbom^{1,*}, Isaac D. Essi¹, Victor A. Kiri²

¹Department of Mathematics, Rivers State University, Port Harcourt, Nigeria

²Department of Mathematics, Physics & Electrical Engineering, Northumbria University, Newcastle upon Tyne, United Kingdom

Abstract Child mortality is a deciding factor of the well-being of a population and often serves as a pointer of the socio-economic status and health development of a nation. A 2017 report on child mortality indicated most child deaths happened in two regions of the world: Sub-Saharan Africa (38%) and South Asia (39%). In Nigeria, around 839,500 children die each year prior to their fifth birthday. This places Nigeria third among the nations of the world with the highest absolute number of child deaths, next to India and Pakistan. Our study is aimed at assessing the effect of socioeconomic, bio-demographic and health-related factors on mortality risk among the under-five (U5M) of Nigerian children, using the 2013 Nigerian Demographic and Health Survey data. We used the non-parametric Kaplan-Meier survival method to assess the influence of the sets of candidate factors on the survival of children aged under-five (U5) and used the Cox proportional hazard model to assess the form of the influence on mortality risk. Those found to have a significant influence on the risk in Nigeria were: mother's educational level, wealth index, marital status, place of residence, sex of the child, region, maternal age at childbirth, number of children ever born, birth interval, and child size at birth. We also found evidence that U5M in Nigeria may be more associated with social, economic, environmental and demographic factors, than with health-related factors. We conclude that for Nigeria to achieve the Sustainable Development Goals, its public health interventions may need to take into account, the factors we have identified.

Keywords Survival analysis, Under-five mortality, Determinants, Nigerian Demographic and Health Survey, Nigeria

1. Introduction

Globally, it was estimated that about 5.4 million children aged under-five die each year, with huge variations and trends across regions and nations [1]. A 2017 UNICEF report on child mortality states that most of these child deaths happened in two particular regions: sub-Saharan Africa (38%) and South Asia (39%). Over half of these deaths occurred in just five nations: the Democratic Republic of the Congo, India, Ethiopia, Nigeria and Pakistan. All things considered, 1 out of every 13 children born in sub-Saharan Africa dies before their fifth birthday as against 1 out of every 185 in high-income countries- meaning that the highest proportions of under-five mortality are concentrated in sub-Saharan Africa and South Asia [1]. About 839,500 Nigeria children die each year before their fifth birthdays.

This places Nigeria third among the countries of the world with the highest absolute number of child deaths, next to India and Pakistan [2-4].

The Millennium Development Goal (MDG) progress report of 2013 revealed that at the national level, only 1/5 of the countries with a high death rate of about 40 deaths per 1000 or more had the chance to achieve MDG Goal 4- the absence of adequate progress in reducing the rates of under-five mortality by many of the developing countries, has resulted in the newly adopted Sustainable Development Goals (SDGs), with the target of reducing U5 mortality to 25 per 1,000 or less by 2030 [5].

While U5 mortality rates have dropped globally by about 56%, that is, from 93 deaths/1,000 live births in 1990 to 41 deaths/1,000 live births in 2016, the rate of progress has been unbalanced across nations with the worst in sub-Saharan Africa where accelerated and sustainable improvements are urgently needed [1]. According to UNICEF 2017, in excess of 50 nations, including Nigeria are missing their SDG targets on child survival, and nearly 60 million children under age 5 will die between 2017 and 2030 if current trends continue.

Nigeria is among the nations in sub-Saharan Africa with

* Corresponding author:

wegbomanthony@gmail.com (Anthony I. Wegbom)

Published online at <http://journal.sapub.org/statistics>

Copyright © 2019 The Author(s). Published by Scientific & Academic Publishing

This work is licensed under the Creative Commons Attribution International

License (CC BY). <http://creativecommons.org/licenses/by/4.0/>

very high under-five mortality [6]. The U5 five mortality rates in 1990 was 201 deaths per 1000, which later dropped to 128 deaths per 1000 in 2013 -marking a 36 per cent drop within the 23-year period. However, this placed Nigeria among the nations that fell far short of achieving the MDG 4 [7]. In any case, there is no evidence that the advancement being made in reducing under-five mortality in Nigeria could be maintained in the post-2015 Millennium Development Goals (MDGs) worldwide plan [8].

In Nigeria, according to the 2013 NDHS report, infant and child mortality is 69 and 64 per 1000 live births respectively. This implies that one in every 15 Nigerian children dies before reaching age 1, and one in every sixteen does not live to their fifth birthday from their first [7]. Factors such as low level of mother's educational level, unsafe sanitation and drinking water, family income, breastfeeding period, birth interval, and place of delivery have been suggested as risks factors of under-five death. This study aims to identify the factors which influence child survival in Nigeria.

2. Materials and Method of Analysis

2.1. Source of Data

Like other surveys that collect data on birth outcomes, the Demographic and Health Surveys (DHS) is a survey that collects data on under-five mortality assessment in developing states like Nigeria, where adequate and reliable vital registration system is missing [9].

The DHS was conducted in 1990, 2003, 2008 and 2013 in Nigeria. This survey is intervallic and cross-sectional directed at the household level. In this study, we used data from the 2013 Nigeria Demographic and Health Survey. The survey is a three-stage stratified cluster design consisting of 904 clusters from which sample of 40,680 households was selected, 45 households in each cluster. A sample of 18,516 women aged 15–49 who had given birth in the past five years preceding the survey was interviewed. Thus, the 2013 NDHS sample was designed to provide estimates for the demographic and health variables at the national level, taking into account, both the rural and urban areas of Nigeria as well as its six geopolitical zones [7].

2.2. Variable Description

The response variable in this study was under-five mortality as a binary with survival status (alive =0 or dead=1) based on age at death in the last five years preceding the survey. In other words, under-five mortality refers to the death of children below the age of five years [7]. The explanatory variables are social, economic, biological, demographic, and health-related variables. The covariates were selected from Mosley and Chen framework [10] as well as some relevant literature and they were accordingly handled as categorical variables: namely, mother educational level (None, primary, secondary, tertiary), wealth index (poor, middle, rich), marital status (never married, married),

religion (Christianity, Islam, others), place of residence (Urban, rural), source of water (unimproved, improved), toilet facility (unimproved, improved), sex of the child (male, female), region (north-central, north-east, north-west, south-east, south-south, south-west), maternal age at childbirth (<25yrs, 25-34yrs, >=35yrs), birth interval (first birth, <24months, 24-35months, >35months), number of children everborn (<=2, 3-5, >6), child size at birth (large, average, small), use bednet (no, yes), place of delivery (public, private, home), and mode of delivery (Cesarean section, other).

2.3. Methods of Analysis

2.3.1. Survival Analysis

We used survival analytical methods, being the conventional approaches for analyzing survival (i.e. time-to-event) data [11-12]. Survival analysis is primarily for studying follow-up times (years, months weeks or days) to the occurrence of any particular event of interest (death or disease) and the technique is particularly very popular in Biostatistics and Demography. Indeed, these methods are the standard statistical techniques of choice for analyzing censored data (i.e. those that containing missing event dates such as where subjects are lost to follow-up) and also for data where the underlying distribution may not be normal [13].

We used the Kaplan-Meier survival technique which is a univariate approach, to identify the effect of each individual variable on the time to death and applied the associated log-rank test to assess the unadjusted effect of each variable on the risk.

In general, the first step in the analysis of survival data is usually to estimate the distribution of the survival times, known as failure times. The survival function is used to describe the lifetimes of the population of interest and subsequently, to estimate the adjusted effects of the explanatory variables.

2.3.2. Formulation

Let the random variable T denote the survival time, and t be the realization of T . Again, let T have a probability distribution function (PDF) of $f(t)$, then the Cumulative distribution function of T can be expressed as

$$F(t) = P(T < t) = \int_0^t f(u)du \quad (1)$$

Survival function $S(t)$ denotes the probability that a child survives longer than t from first day of birth and is expressed as:

$$S(t) = P(T > t) \quad (2)$$

$$S(t) = 1 - P(\text{a child survives } > t \text{ from first day of birth}) \quad (3)$$

Hazard function, $h(t)$ represents the probability that a child dies at time t , on the condition that he survived to that time. It is the instantaneous death rate at which a child survives to time t :

$$h(t) = \lim_{\Delta t \rightarrow 0} \frac{P(t < T \leq t + \Delta t | T > t)}{\Delta t} = \frac{f(t)}{S(t)} \quad (4)$$

Where Δt is the instantaneous change.

The survival analysis model used to obtain the adjusted effects of the variables in our study was the Cox proportional hazard model (PHM). The PHM is a useful survival analytical technique for accessing the influence of relevant factors on the hazard function. The continuous random variable denotes the lifetime t of a child and X as the vector of explanatory factors. The PHM

$$h_i(t, X) = h_0(t)\exp(\beta_1X_1 + \beta_2X_2 + \beta_3X_3 + \dots + \beta_pX_p) \quad (5)$$

$$h_i(t, X, \beta) = h_0(t)\exp\sum_{i=1}^p \beta X_i \quad (6)$$

Where

$h_i(t, X)$ is the risk of death for child i at time t ; $h_0(t)$ is the baseline hazard function; $X = (x_1, x_2, x_3, \dots, x_p)$, are the selected variables; β is the vector of unknown coefficients of the explanatory variables x_i .

The key assumption of the Cox regression model is that the risk for child i , is proportional to the risk for any other child j . That is, the ratio of their risks (i.e. the hazard ratio) is constant over time t .

If the vector $X_i = 0$, then the hazard function for the i th individual becomes the baseline hazard function. However, for any other value of X_i , we can divide both sides of equation (5) above by $h_0(t)$, to obtain

$$\frac{h_i(t)}{h_0(t)} = \exp\sum_{i=1}^p \beta_i X_i \quad (7)$$

The hazard ratio (HR) represents the instantaneous risk over the study time period and as a risk estimator, it is generally more robust to selection bias than other risk estimators such as Odds ratio and relative risk. It can be interpreted as the chance of an event occurring in one group/category of a factor, divided by the chance of the event occurring in the other group/category. Conventionally, a 95% confidence interval (CI) for the HR is usually provided for each variable and its effect on the risk is considered as statistically significant if the CI does not include the value of 1 [13].

All analyses were conducted using the STATA statistical package (version 15).

3. Results

3.1. Univariate Survival Statistics

Table 1a and Table 1b show those among the list of bio-demographic and health-related variables and the socio-economic variables respectively, on which we found evidence of association with the risk of under-five mortality based on the log-rank test (i.e. $p < 0.05$). The two sets of results suggest the following variables as possibly associated with the risk of death before the age of five: namely, the sex and size of the child, geopolitical region, the maternal age of the mother at birth, the birth interval, number of children ever-born, usage of bednet, place and mode of delivery, the educational level of the mother, wealth index, marital status,

religion, place of residence, source of water, and toilet facility as possibly associated with the risk of death before the age of five.

Figure 1 shows the overall Kaplan Meier estimate of the survival function. The result shows that most children died in the early months of birth, with the risk gradually declining as they age but which becomes less noticeable as they approach the age of five.

Table 1a. Log-rank test for equality of survival times between the different categories for each of the bio-demographic and health-related characteristics

Background characteristics	Death among under-5 children (%)	Chi-square (DF)	p-value
Sex of the child		15.12(1)	0.0001
Male	8.4		
Female	7.1		
Region		2589.04(5)	0.0000
North central	1.3		
North east	3.6		
North west	7.0		
South east	1.2		
South south	1.2		
South west	1.2		
Maternal age at childbirth		187.03(2)	0.0000
<25yrs	9.0		
25-34yrs	5.3		
>=35yrs	1.3		
Birth interval		1281.60(3)	0.0012
First birth	3.6		
<24months	5.6		
20-35moths	4.2		
>35months	2.1		
Number of children ever born		1734.11(2)	0.0000
<=2	0.8		
3-5	3.8		
6+	10.9		
child size at birth		36.87(2)	0.0000
Large	6.9		
Average	6.3		
Small	2.3		
Use bednet		9.92(1)	0.0016
No	12.5		
Yes	3.1		
Place of delivery		137.30(2)	0.0000
Public	3.5		
Private	2.4		
Home	9.6		
Mode of delivery		12.42(1)	0.0004
Cesarean section	3.1		
Others	12.4		

Table 1b. Log-rank test for equality of survival times between the different categories for each of the socio-economic variables


Background characteristics	Death among under-5 children (%)	Chi-square	P-value
Educational level		2054.29	0.0000
None	10.2		
Primary	3.0		
Secondary	1.9		
Tertiary	0.4		
Wealth index		2439.23	0.0000
Poor	9.8		
Middle	2.7		
Rich	3.0		
Marital status		12.34	0.0004
Never married	1.0		
Married	14.5		
Religion		1390.75	0.0000
Christianity	4.5		
Islam	10.8		
Others	0.3		
Place of residence		796.96	0.0001
Urban	3.6		
Rural	11.9		
Source of water		114.89	0.0000
Unimproved	7.7		
Improved	7.9		
Toilet facility		133.09	0.0021
Unimproved	8.7		
Improved	6.8		

3.2. Risks Factors of Under-Five Mortality

Table 2 shows the risk factors associated with under-five mortality in Nigeria based on three forms of the Cox proportional hazards model. Model 1 involved only the socio-economic variables, whereas model 2 additionally involved the bio-demographic variables but without the health-related one. Finally, model 3 included all three sets of variables, namely the socio-economic, bio-demographic and

health-related types. The results indicate the following as significantly contributing to the risk of under-five mortality in Nigeria: namely, the educational level of the mother, wealth index, marital status, Place of residence, Sex of the child, Geographical Region, Maternal age at childbirth, Birth interval, Number of children ever-born, and the Size of the child at birth.

More specifically, the higher the educational level of the mother, the lower the risk of under-five death. Children whose mothers had tertiary and secondary levels of education were less likely to die compared to children whose mother had none education (HR = 0.67, 95% CI = 0.46-0.98) and (HR = 0.78, 95% CI = 0.64-0.96) respectively. Households with rich and middle wealth index also had a lower risk of dying before the age of five, compared to households with poor wealth index (HR = 0.71, 95% CI = 0.59-0.87) and (HR = 0.73, 95% CI = 0.63-0.85) respectively. Children from married parents were less likely to die within five years of existence (HR = 0.71, 95% CI = 0.57-0.91) than children whose parents were never married. The likelihood of under-five mortality was 1.29 (95% CI = 1.11–1.50) times higher among children residing in rural areas when compared with their urban counterpart.

**Figure 1.** Plot of the overall Kaplan-Meier estimate of the survivor function for the under-five children in Nigeria**Table 2.** Cox's proportional hazards model: risk factors of under-five mortality in Nigeria

Covariates	Model 1 HR (95% CI)	Model 2 HR (95% CI)	Model 3 HR (95% CI)
Educational level			
None	1	1	1
Primary	0.82 (0.78-0.87)*	0.92 (0.79-1.08)	0.94 (0.80-1.09)
Secondary	0.69 (0.64-0.74)*	0.77 (0.63-0.94)*	0.78 (0.64-0.96)*
Tertiary	0.49 (0.43-0.56)*	0.64 (0.44-0.92)*	0.67 (0.46-0.98)*
Wealth index			
Poor	1	1	1
Middle	0.67 (0.63-0.70)*	0.72 (0.62-0.84)*	0.73 (0.63-0.85)*
Rich	0.51 (0.48-0.55)*	0.69 (0.57-0.84)*	0.71 (0.59-0.87)*
Marital status			
Never married	1	1	1

Married	0.84 (0.78-0.90)*	0.72 (0.57-0.91)*	0.71 (0.57-0.91)*
Religion			
Christianity	1	1	1
Islam	1.35 (1.28-1.42)	1.10 (0.92-1.33)	1.12 (0.93-1.35)
Others	1.04 (0.90-1.19)	0.71 (0.44-1.13)	0.68 (0.42-1.10)
Place of residence			
Urban	1	1	1
Rural	1.09 (1.04-1.14)*	1.32 (1.14-1.54)*	1.29 (1.11-1.50)*
Source of water			
Unimproved	1	1	1
Improved	1.03 (0.99-1.07)	0.97 (0.88-1.08)	0.97 (0.87-1.08)
Toilet facility			
Unimproved	1	1	1
Improved	1.02 (0.98-1.06)	0.95 (0.85-1.06)	0.95 (0.85-1.06)
Sex of the child			
Male		1	1
Female		0.91 (0.82-1.01)	0.91 (0.82-1.00)*
Region			
North central		1	1
North east		1.14 (0.93-1.39)	1.13 (0.92-1.39)
North west		1.41 (1.16-1.72)*	1.39 (1.12-1.70)*
South east		1.88 (1.46-2.42)*	2.04 (1.57-2.65)*
South south		1.10 (0.85-1.42)	1.09 (0.84-1.41)
South west		1.01 (0.77-1.31)	1.06 (0.82-1.39)
Maternal age at child birth			
<25yrs		1	1
25-34yrs		0.82 (0.72-0.94)*	0.82 (0.72-0.94)*
>=35yrs		0.80 (0.66-0.97)*	0.80 (0.66-0.97)*
Birth interval			
First birth		1	1
<24months		0.91 (0.73-1.13)	0.92 (0.74-1.15)
24-35moths		0.66 (0.53-0.83)*	0.68 (0.54-0.85)*
>35months		0.49 (0.39-0.62)*	0.50 (0.39-0.63)*
Number of children everborn			
<=2		1	1
3-5		1.45 (1.18-1.78)*	1.41 (1.15-1.73)*
6+		2.12 (1.68-2.68)*	2.08 (1.65-2.63)*
Child size at birth			
Large		1	1
Average		1.10 (0.99-1.23)	1.09 (0.97-1.22)
Small		1.30 (1.13-1.49)*	1.28 (1.11-1.48)*
Use bednet			
No			1
Yes			0.96 (0.84-1.08)
Place of delivery			
Public			1
Private			0.93 (0.74-1.18)
Home			1.11 (0.95-1.31)
Mode of delivery			
Cesarean section			0.77 (0.44-1.34)
Others			1

Female children were less likely to die on or before reaching age five compared to male children (HR = 0.91, 95% CI = 0.82–1.00). Residing in the north-west and south-east geopolitical regions was significantly associated higher likelihood of under-five death (HR = 1.39, 95% CI = 1.12–1.70) and (HR = 2.04, 95% CI = 1.57–2.65) respectively when compared to the north-central region. Children whose mothers were aged under 25 at the time of giving birth had the highest risk of dying before the age of five: (HR = 0.82, 95% CI = 0.72–0.94) and (HR = 0.80, 95% CI = 0.66–0.97) for mothers aged 25–34 years and ≥ 35 years respectively. There was a declining probability of mortality for children with large and medium birth interval (HR = 0.50, 95% CI = 0.39–0.63) and (HR = 0.68, 95% CI = 0.54–0.85) respectively when compared to those with short birth interval. The number of children ever-born was associated with a higher risk of U5M. Children born in a family of 3–5 and 6 or more were 1.41 and 2.08 times more likely to die than children born in a family with ≤ 2 number of children ever born. The size of the child at birth also had an influence on the risk of death. Children with Small size at birth had a higher risk of death (HR = 1.28, 95% CI = 1.11–1.48) compared with large size.

4. Discussion

In recent years, the under-five mortality rates have been declining consistently in Nigeria which indicates a level of progress in the quality of life. These rates are additionally vital to public health programmes in Nigeria [1, 7, 10]. The results of this study suggest that the risk of Under-5 death depends greatly on a number of socio-economic, bio-demographic and health-related variables, such as mother educational level, wealth index, marital status, place of residence, gender, geopolitical region, mother's age at birth, preceding birth interval, number of children ever born, and the size of child at birth.

Mother's education had a significant relationship with child survival. Higher risks of children death were identified among the illiterate mothers compared with literate mothers: mothers with secondary or higher education mothers. This is consistent with the findings by others [13–17]. A possible explanation could be that educated mothers have better socioeconomic status, better knowledge on family health and childcare, preventive care, effective use of modern health services and good management of child illness [18]. According to O'Toole, education additionally changes the customary and social familial relationships as regards decision making and engages the mothers in several issues like childcare which plays a role in reducing child mortality [19].

Substantial disparities in mortality risks were also observed by maternal age at the birth of the child. We found evidence of a higher risk of death for children of younger mothers and this also confirms the findings by previous research [20–23]. Mother's age at the birth of a child can influence child mortality through different standpoints: the

higher risk of child death among younger mothers may be attributed to immature reproductive systems and less stability to handle the complexities of pregnancy [14]. Furthermore, we found the marital status of the mother as related to the risk of child death. There is a low probability of children of married mothers dying compared to those born to never-married mothers- this was also the finding by an earlier study by one of the authors in rural Nigeria [21].

The risk of under-5 mortality was similarly lower for female children. This is in line with the findings of other studies which have suggested girls as more biologically advantaged than boys in terms of many of the causes of death [24–25, 17]. We also found evidence that the length of preceding birth interval and the number of children ever-born were associated with the risk of death before the age of five. On birth intervals, we found evidence that children delivered within shorter birth intervals were at a greater risk of dying before their fifth birthday. Thus, agreeing with the findings by some previous studies [26–29, 16, 21, 23, 14, 17]. A possible explanation is that shorter length of the birth interval may negatively influence maternal health, by increasing susceptibility of infectious diseases and competition in familial resource among children [27].

Our findings showed that survival of children to their fifth year in life was significantly worse among children residing in the north-west and south-east parts of Nigeria than those residing in the North central and other part zones. This high risk might be attributed to social improvement in the community, population density, territorial advancement, as well as regional economic resources [17, 30–32]. We also found evidence that the place of residence of the mother and her wealth index might influence the survival of her child before the fifth birthday. On the wealth status of the household, we also found evidence that mortality risk in this age group might be lower in wealthy households as compared to the underprivileged households. This was consistent with the findings by some previous studies [33–35]. These households have better housing conditions, better nutrition, more empowerment and hence they may be able to afford better medical attention and care thus significantly enhancing the survival probability of all their children [33].

Our study also found evidence that the child's size at birth might be significantly associated with infant mortality. Although the size at birth may not be so important once a child survives beyond the first year of life, timely intervention in the form of promoting breastfeeding and micronutrient interventions is recommended for stunted and underweight children, particularly during the neonatal period and late infancy [36–38]. Indeed, Uthman had noted that underweight children are more vulnerable to infections and other related diseases [39].

The size of the child at birth was found to be associated with high U5M and the evidence also supports the findings by others that small sized children at birth have a higher risk of dying before the age of five [17, 36–39]. This might be due

to the immaturity of their organs which makes them find it more difficult to adapt to eternal, independent life. Most premature babies are prone to have sepsis which is one of the major causes of Under 5 Mortality [30].

Furthermore, our study suggests that mortality rates are higher in rural areas. This concurs with previous studies in and outside Nigeria [40-43, 30, 17]. This is on the ground that those living in the urban areas have access to improved water supply, improve sanitation facilities, unlimited access to healthcare as well as other social and economic services [30, 43].

5. Conclusions

The findings from this study have essential policy implications, particularly in monitoring public health interventions which need to ensure a consistent decline in child mortality rates, towards the achievement of the SDG 3. In this study, we explored and then identified the socioeconomic, bio-demographic, and health-related factors that might influence the risk of U5M in Nigeria, using the method of time-to-event (i.e. survival) analysis. The results from the Cox regression models showed that mother educational level, wealth index, marital status, place of residence, gender, geopolitical region, mother's age at birth, preceding birth interval, number of children ever born, and the size of the child at birth might influence the risk of child mortality in Nigeria. In a nutshell, these findings suggest that under-five mortality is greatly associated with socio-economic and bio-demographic factors in addition to those already known as health-related.

REFERENCES

- [1] UNICEF (2017). Levels and Trends in Child Mortality: Report 2017. Estimates Developed by the UN Inter-agency Group for Child Mortality Estimation. Available from: http://www.childmortality.org/files_v20/download/igme%20report%202015%20c.
- [2] World factbook (2018). <https://www.cia.gov/library/publications/the-world-factbook/rankorder/2091rank.html>.
- [3] National Bureau of Statistics (NBS) and United Nations Children's Fund (UNICEF) (2017). 2017 Multiple Indicator Cluster Survey 2016-17, Survey Findings Report. Abuja, Nigeria.
- [4] WHO, (2019). www.premiumtimesng.com/news/headlines/246720-nigeria-third-highest-infant-mortality-rate-world.html
- [5] United Nations (2015). Transforming our World: the 2030 Agenda for Sustainable Development. United Nations. Retrieved April 28, 2018, from http://www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E.
- [6] UNICEF (2014). Levels and Trends in Child Mortality: Report 2014. Estimates Developed by the UN Inter-agency Group for Child Mortality Estimation.
- [7] National Population Commission (NPC) [Nigeria] and ICF International (2014). Nigeria Demographic and Health Survey 2013. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF International.
- [8] National Bureau of Statistics [Nigeria] (2014). The Millennium Development Goals Performance Tracking Survey 2015 Report. 2014 MDG Survey Report, Abuja. Retrieved April 28th, 2018, from <http://www.nigerianstat.gov.ng/report/254>.
- [9] Pedersen J, Liu J (2012). Child Mortality Estimation: Appropriate Time Periods for Child Mortality Estimates from Full Birth Histories. *PLoS Med.* 9(8): e1001289. doi:10.1371/journal.pmed.1001289.
- [10] Mosley, WH and Chen LC (1984). An Analytical Framework for the Study of Child Survival in Developing Countries. *Population and Development Review*, 10, 25- 45. Retrieved from <http://www.jstor.org/stable/2807954>.
- [11] Lee TE and Wang JW (2003). *Statistical Methods for Survival Data Analysis*. 3rd ed. A John Wiley and Sons, Inc., Publication.
- [12] David Collett (2003). *Modelling Survival Data in Medical Research*. Text in Statistical Sci.2nd ed. Chapman & Hall/CRC.
- [13] Kleinbaum D and Mitchel K (2012). *Survival Analysis: A Self-Learning Text*.3rd ed. Springer Science + Business Media, Inc.
- [14] Khan and Awan (2017). A comprehensive analysis on child mortality and its determinants in Bangladesh using frailty models *Archives of Public Health*. 75: 58. DOI 10.1186/s13690-017-0224-6.
- [15] Fayehun OA (2010). Household Environmental Health Hazards and Child Survival in sub-Saharan Africa. DHS Working Papers No. 74. Calverton, Maryland, USA: ICF Macro.
- [16] Wegbom AI, Wokoma DSA, Nnoka LC, Chibo O (2016a). What explains the high rate of infant mortality in rural Nigeria: biodemographic or socioeconomic factors? *Int J Health Sci Res.* 6(8): 317-323.
- [17] Adebowale SA, Morakinyo OM and Ana GR (2017). Housing materials as predictors of under-five mortality in Nigeria: evidence from 2013 demographic and health survey *BMC Pediatrics* 17:30 DOI 10.1186/s12887-016-0742-3.
- [18] Caldwell JC (1994). How is greater maternal education translated into lower child mortality? *Health Transit Rev.* 4(2): 224-9.
- [19] O'Toole J, Wright RE (1991). Parental education and child mortality in burundi. *J Biosoc Sci.* 23(3): 255-62.
- [20] Akinyemi JO, Bamgboye EA, Ayeni O (2015). Trends in neonatal mortality in Nigeria and effects of bio-demographic and maternal characteristics. *BMC Paediatr.* 15:36. doi:10.1186/s12887-015-0349-0.
- [21] Wegbom AI, Aboko IS, Mandah OCA (2016b). Multivariate Analysis of Child Mortality determinants in Rural Nigeria. *Tlep Int. Journal of Multi Discipline in Industrial and Applied Sciences.* 1(8): 1-10.
- [22] Alam N (2000). Teenage motherhood and infant mortality in

- Bangladesh: Maternal age [hyphen] dependent effect of parity one. *J Biosoc Sci.* 32(02): 229–36.
- [23] Getachew Y, Bekele S (2016). Survival Analysis of Under-Five Mortality of Children and its Associated Risk Factors in Ethiopia. *J Biosens Bioelectron* 7: 213. doi: 10.4172/2155-6210.1000213.
- [24] Alonso V, Fuster V, Luna F (2006). Causes of neonatal mortality in Spain (1975–98): influence of sex, rural-urban residence and age at death. *J Biosoc Sci.* 38(04): 537–51.
- [25] Sawyer CC (2012). Child mortality estimation: estimating sex differences in childhood mortality since the 1970s. *PLoS Med.* 9(8): 1001287.
- [26] Akinyemi JO, Bamgboye EA and Ayeni O (2013). New trends in underfive mortality determinants and their effects on child survival in Nigeria: A review of childhood mortality data from 1990-2008; *African Population Studies.* 27.1: 25-42.
- [27] Rutstein SO (2005). Effects of preceding birth intervals on neonatal, infant and under-five years mortality and nutritional status in developing countries: evidence from the demographic and health surveys. *Int J Gynecol Obstet.* 89:7–24. 49.
- [28] Whitworth A, Stephenson R (2002). Birth spacing, sibling rivalry and child mortality in india. *Soc Sci Med.* 55(12): 2107–19. 50.
- [29] Mustafa HE, Odimegwu C (2008). Socioeconomic determinants of infant mortality in kenya: analysis of kenya dhs 2003. *J Humanit Soc Sci.* 2(8): 1934–722.
- [30] Kayode GA, Adekanmbi VT and Uthman OA (2012). Risk Factors and a predictive model for underfive mortality in Nigeria: evidence from Nigeria demographic and health survey. *BMC Pregnancy and Childbirth.*
- [31] Adebayo SB and Fahrmeir L (2002). Analyzing Child Mortality in Nigeria with Geoadditive Survival Models. *Sonderforschungsbereich 386, Paper 303.*
- [32] Antai D (2011). Regional Inequalities in Under-five Mortality in Nigeria: A Population-based Analysis of Individual - and -Community Determinants. *Pop. Health Metrics,* 9(6): 1-10 Available from <http://www.pophealthmetrics.com/content/9/1/6>.
- [33] Mutunga CJ (2007). Environmental determinants of child mortality in Kenya. United Nations University. Research Paper No. 2007/83.
- [34] Uddin J, Hossain Z and Ullah MO (2009). Child Mortality in a Developing Country: A Statistical Analysis. *Journal of Applied Quantitative Methods,* 4(3): 270-283.
- [35] Bello RA and Joseph AI (2014). Determinants of Child Mortality in Oyo State, Nigeria. *African Research Review,* 8(1): 252-272. DOI: <http://dx.doi.org/10.4314/afrr.v8i1.17>.
- [36] Bhutta Z, Bawany FA, Feroze, AB, Rizvi A, Thapa SJ. and Patel M (2009). Effects of the Crises on Child Nutrition and Health in East Asia and the Pacific. *Global Social Policy,* 9(1 suppl), 119- 143. doi: 10.1177/1468018109106888.
- [37] Black RE, Allen LH, Bhutta ZA, Caulfield LE, de Onis M, Ezzati M and Rivera J (2008). Maternal and child undernutrition: global and regional exposures and health consequences. *The Lancet,* 371(9608), 243-260. doi: 10.1016/s0140-6736(07)61690-0.
- [38] Imdad A, Yakoob M and Bhutta ZA (2011). Impact of maternal education about complementary feeding and provision of complementary foods on child growth in developing countries. *BMC Public Health,* 11(3). doi: 1471-2458/11/S3/S25.
- [39] Uthman OA (2008). Effect of low birth weight on infant mortality: Analysis Using Weibull Hazard Model. *The Internet Journal of Epidemiology.* 6 (1). DOI: 10.5580/fca.
- [40] Ezeh OK, Agho KE, Dibley MJ, Hall JJ, Page AN (2015). Risk factors for postneonatal, infant, child and under5 mortality in Nigeria: a pooled cross-sectional analysis. *BMJ Open [Internet].* 5:e006779.
- [41] Kazembe L, Clarke A, Kandala NB (2012). Childhood mortality in sub-Saharan Africa: cross-sectional insight into small-scale geographical inequalities from Census data. *BMJ Open.* 2:1–12.
- [42] Morakinyo OM, Fagbamigbe AF (2017). Neonatal, infant and under-five mortalities in Nigeria: An examination of trends and drivers (2003-2013). *PLoS ONE* 12(8): e0182990. <https://doi.org/10.1371/journal.pone.0182990>.
- [43] Dejene T, Girma E (2013). Social determinants of under-five mortality in Ethiopia: Event history analysis using evidence from Ethiopian Demographic and Health Survey (EDHS). *Health (Irvine. Calif);* 5:879– 84.